

<b>(51) International Patent Classification 6 :</b> <b>C12P 17/02, C12N 3/00, 1/00, 15/01, 1/20, C12P 7/62, 7/26 // (C12P 17/02, C12R 1:07) (C12P 17/02, C12R 1:125)</b>		<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/32651</b> <b>(43) International Publication Date:</b> 1 July 1999 (01.07.99)
<b>(21) International Application Number:</b> PCT/CA98/01150 <b>(22) International Filing Date:</b> 18 December 1998 (18.12.98) <b>(30) Priority Data:</b> 08/995,960 22 December 1997 (22.12.97) US <b>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application</b> US 08/995,960 (CIP) Filed on 22 December 1997 (22.12.97) <b>(71) Applicant (for all designated States except US):</b> BCM DEVELOPEMENT INC. [CA/CA]; Suite 218, 125 rue Dalhousie, Québec, Québec G1K 4C5 (CA). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> PAGE, Michel [CA/CA]; Suite 218, 125 rue Dalhousie, Québec, Québec G1K 4C5 (CA). LANDRY, Nathalie [CA/CA]; 1060 de la Voie Ouest, St-Jean-Chrysostome, Québec G6Z 1K1 (CA). BOISSINOT, Maurice [CA/CA]; 109 Jean Bruchési, Saint-Augustin, Québec G3A 2N2 (CA). HELIE, Marie-Claude [CA/CA]; 453 Place St-Laurent,		Cap-Rouge, Québec G1Y 3G9 (CA). HARVEY, Mario [CA/CA]; 1067-F De la Prairie, St-Jean-Chrysostome, Québec G6Z 2G3 (CA). GAGNE, Martin [CA/CA]; 6400 Pise, Charlesbourg, Québec G1H 7A7 (CA). <b>(74) Agents:</b> COTE, France et al.; Swabey Ogilvy Renault, Suite 1600, 1981 McGill College Avenue, Montréal, Québec H3A 2Y3 (CA). <b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> BACTERIAL MASS PRODUCTION OF TAXANES AND PACLITAXEL			
<b>(57) Abstract</b> <p>The present invention relates to a method of obtaining different biologically pure cultures of bacteria isolated from different species of <i>Taxus</i> such as <i>Taxus canadensis</i>, <i>T. brevifolia</i>, <i>T. baccata</i>, <i>T. cuspidata</i>, and <i>T. hunnewelliana</i>, wherein the bacteria produce <i>in vitro</i> taxanes and paclitaxel, and wherein the bacteria are of the genus <i>Sphingomonas</i>, <i>Bacillus</i>, <i>Pantoea</i> or <i>Curtobacterium</i>. Also, the present invention relates to a method of a bacterial mass production of at least one taxane or paclitaxel. There is also disclosed a novel bacterial taxane. The present invention also relates to the use of different biologically pure cultures of bacteria isolated from different species of <i>Taxus</i>, wherein the bacteria are able to biotransform pro-taxanes. There is also provided a process for improving taxanes and paclitaxel production of taxanes and paclitaxel producing bacteria which include culturing bacteria in the presence of a mutagenic agent for a period a time sufficient to allow mutagenesis. There is disclosed two new mutated bacterial which have an increased yield of pro-taxane biotransformation.</p>			

**BEST AVAILABLE COPY**

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Licchtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

BACTERIAL MASS PRODUCTION OF TAXANES AND PACLITAXELBACKGROUND OF THE INVENTION(a) Field of the invention

5           The present invention relates to the production of paclitaxel and derivatives thereof (such as related taxanes) using a plurality of different bacteria isolated from different species of *Taxus*, and also to a novel taxane. There are disclosed methods for the isolation of these bacteria and the screening tests that  
10           were used to provide evidence that paclitaxel and taxanes were produced by said bacteria. There are also disclosed methods for the biotransformation of pro-taxanes by said bacteria.

15   (b) Description of prior art

          Paclitaxel, also referred to as Taxol™, has been first identified in 1971 by Wani and collaborators (Wani MC et al., 1971 *J. Am. Chem. Soc.*, 93: 2325-2327) following a screening program of plant extracts of the  
20   National Cancer Institute. This complex diterpene showed cytotoxic activity against several types of tumors and is presently used in the treatment of some cancers such as ovarian and breast cancers. Clinical studies suggest that Taxol™ could eventually be used in  
25   the treatment of over 70% of human cancers.

          Paclitaxel differs from other cytotoxic drugs by its unique mechanism of action. It interferes with cell division by manipulating the molecular regulation of the cell cycle. Paclitaxel binds to tubulin, the  
30   major structural component of microtubules that are present in all eukaryotic cells. Unlike other antimitotic agents such as vinca alkaloids and colchicine,

which inhibit the polymerization of tubulin, paclitaxel promotes this assembly of tubulin and stabilizes the resulting microtubules. This event leads to the interruption of cell division, and ultimately to cell death.

5           The major obstacle in the use of paclitaxel as an anticancer treatment is its supply. It was originally isolated from the bark and leaves of yew trees such as *Taxus brevifolia*, *T. baccata*, *T. cuspidata* or, *T. canadensis*. The low yield of the isolation of  
10 paclitaxel (0,016 g%) and the limited availability of the trees have forced the scientific and industrial community to find alternative ways of producing paclitaxel.

          The antitumor property of taxoid compounds has  
15 also lead to the generation of new anticancer drugs derived from taxanes. Taxotere™ (sold by Rhône-Poulenc Rorer), which is produced from 10-deacetylbaccatin III by hemisynthesis, is currently used in the treatment of ovarian and breast cancers. Furthermore, Abbott Laboratories  
20 is conducting clinical trials with a drug derived from 9-dihydro-13-acetyl baccatin III, a natural precursor specific to *Taxus canadensis*. The increasing demand for taxanes highlights the urgent need for renewable and economical processes that would  
25 not endanger plant species.

          Presently, industrials are producing paclitaxel through hemisynthesis from baccatin III, a natural precursor of paclitaxel. However, this process still relies on a plant substance that must be extracted from  
30 yew trees. The first complete chemical synthesis of paclitaxel has been achieved in 1994 by Nicolaou et al. (1994, *Nature*, 367:630-634). This is a multistep proc-

ess and the overall yield has made this approach non economically feasible.

Plant cell culture of *Taxus* species is another approach explored by many groups (Yukimune et al., 1996, *Nature Biotechnology*, 14:1129-1132; Srinivasan et al., 1995, *Biotechnology and Bioengineering*, 47:666-676). Somehow, this process is limited by the amount of paclitaxel that can be produced, the length of incubation time required to obtain significant yields, and the application of plant cell culture to the large volumes required by the industry.

In United States Patent No. 5,322,779, in the names of Gary A. Strobel et al. disclosed a fungus isolated from the bark of a sample of *Taxus brevifolia* which is able to synthesize paclitaxel at a level of 24-50 ng/l after a period of incubation of 3 weeks. Later, Strobel et al. (1996, *Microbiology*, 142:435-440) reported another fungus, *Pestalotiopsis microspora*, isolated from the inner bark of *Taxus wallachiana* that can produced up to 55 µg/l of paclitaxel within 5 weeks. Somehow, the long periods of incubation and the large volumes required to extract significant amounts of paclitaxel reduce the profitability of the process.

In United States Patent number 5,561,055 (issued on October 1, 1996 in the names of Michel Pagé et al., the Applicant), there is disclosed one bacterium, which was referred to as *Erwinia taxi*, for the production of paclitaxel. Since then, this bacteria has been characterized as *Sphingomonas taxi*. This bacterium was isolated from *Taxus canadensis*. It would be highly desirable to be provided with other bacteria having

highly diverse metabolic capacities isolated from different species of *Taxus* for the production of paclitaxel and related taxanes at higher yields.

It would also be highly desirable to be provided with widely different bacteria for the mass production of various different bacterial taxanes.

As mentioned in International Patent Application published under number WO97/16200, biotransformation process may be used for the generation of new taxanes molecules that lead to new therapeutic drugs. It would also be highly desirable to be provided with new strains of microorganisms able to biotransform taxanes compounds for use as therapeutic agents or to be modified by hemisynthesis.

Genetic manipulations of bacteria can increase the activity or the production of certain proteins. It would also be highly desirable to be provided with mutant of our original isolates that could produce and biotransform taxanes at higher levels.

#### SUMMARY OF THE INVENTION

One aim of the present invention is to provide a plurality of bacteria for the mass production of taxanes and paclitaxel.

Another aim of the present invention is to provide a method for bacterial mass production of taxanes and paclitaxel which overcomes all the drawbacks of the prior art.

Another aim of the present invention is to provide a novel process for the production of taxanes and paclitaxel. The industrial application of this process would provide alternative renewable sources of taxoids



compounds for the pharmaceutical industry.

Another aim of the present invention is to provide a biotransformation process in which plant-derived taxanes are converted into substances that may be useful for the production of other therapeutic compounds.

In accordance with the present invention there is provided a method to obtain biologically pure cultures of bacteria isolated from *Taxus*, wherein said bacteria produce de novo taxanes and paclitaxel at a concentration of about 1 to 25 µg/L, wherein said bacteria are isolated from the inner surfaces of different species of *Taxus* including without limitations *Taxus canadensis*, *T. brevifolia*, *T. hunnewelliana*, *T. baccata*, and *T. cuspidata*.

In addition, said bacteria are capable of producing biotransformed taxanes wherein pro-taxanes are added to their culture medium.

Such biologically pure cultures of bacteria of the present invention include, without limitation, bacteria of the genus selected from the group consisting of *Sphingomonas*, *Bacillus*, *Pantoea*, and *Curtobacterium*.

In accordance with the present invention, the bacteria include, without limitation, *Bacillus cereus* ssp. *taxi*, *Bacillus megaterium* ssp. *taxi*, *Pantoea* sp. BCM 1, *Pantoea* sp. BCM 2, *Pantoea* sp. BCM 3, *Bacillus cereus* ssp. BCM 4, *Bacillus subtilis* ssp. *taxi*, *Bacillus megaterium* ssp. BCM 9 or *Curtobacterium* sp. BCM 5.

In accordance with the present invention there is also provided a method of bacterial mass production of taxanes and paclitaxel thereof which comprises the steps of:

- 5 a) culturing the bacteria of the present invention in a growth-supporting nutrient medium capable of promoting growth and reproduction of said bacteria, and wherein said culturing is effected for a time sufficient to allow production of taxanes and paclitaxel; and
- b) isolating said produced taxanes and paclitaxel from said bacteria or culturing medium of step a).

10 In accordance with the present invention there is also provided a process for improving taxanes and paclitaxel production of taxanes and paclitaxel producing bacteria comprising the steps of:

- 15 a) culturing bacteria in the presence of a mutagenic agent for a period of time sufficient to allow mutagenesis;
- b) selecting said mutants by a change of the phenotype which results in an increased production of taxanes and paclitaxel.

20 The mutagenic agent may be a chemical agent, such as daunorubicin and nitrosoguanidine.

The mutagenic agent may be a physical agent, such as gamma radiation or ultraviolet.

25 The mutagenic agent may be a biological agent, such as a transposon.

In accordance with the present invention, the mutated bacteria include, without limitation, *Sphingomonas taxi* D200 or *Sphingomonas taxi* D201.

30 In accordance with the present invention there is also provided a method of bacterial biotransformation of taxanes and paclitaxel thereof which comprises the steps of:



5 a) incubating the bacteria of the present invention in a growth-supporting nutrient medium capable of promoting growth and reproduction of said bacteria, and wherein said incubation is effected in the presence of pro-taxanes for a time sufficient to allow production of taxanes and paclitaxel; and

b) isolating said produced taxanes and paclitaxel thereof from said culturing medium of step a).

10 In accordance with the present invention, there is also provided a process for improving biotransformation of pro-taxanes into taxanes and paclitaxel by taxanes and paclitaxel-producing bacteria comprising the steps of:

15 a) culturing bacteria in the presence of a mutagenic agent for a time sufficient to allow mutagenesis; and

20 b) selecting said mutants by a change of the phenotype which results in an increased biotransformation of pro-taxanes into taxanes and paclitaxel.

In accordance with the present invention there is also provided a method of bacterial biotransformation of pro-taxanes into taxanes and paclitaxel thereof which comprises the steps of:

25 a) incubating the mutated bacteria of the present invention in a nutrient medium, and wherein said incubation is effected in the presence of pro-taxanes for a time sufficient to allow production of taxanes and paclitaxel; and

30 b) isolating said produced taxanes and paclitaxel thereof from said culturing medium of step a).


In accordance with the present invention there is also provided a novel bacterial taxane having a characteristic ultraviolet spectrum in HPLC as shown in Fig. 9B, Electrospray Ionization (EI) spectrum as shown in Fig. 10A, and characteristic fragments as shown in Fig. 10B.

This novel bacteria-derived taxane of the present invention is produced by at least two species of *Bacillus* (*B. cereus* ssp. *taxi* and *B. megaterium* ssp. *taxi*).

For the purpose of the present invention the following terms are defined below.

The term "taxanes and paclitaxel" is intended to mean any paclitaxel derivatives or precursor which have retained or not the taxol-associated cytotoxic biological activity or are thought to be precursors in the synthesis of paclitaxel. Such taxanes and paclitaxel may be selected from the group consisting of all the diterpenes isolated from any *Taxus* species. The production of all taxanes by bacteria, whether pharmacologically active or inactive, is contemplated within the scope of the present invention. Taxanes that are produced by the bacteria of the present invention may be as such found in *Taxus* plant species or may differ from the ones found in *Taxus* plant species.

Exemplary taxanes which may be produced by the bacteria of the present invention include but are not limited to those of the following Formula I:



I

wherein  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  
5  $R_{13}$ ,  $R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$  are defined in Table 1 below.

However, since bacteria have highly diverse metabolic capacities the taxanes and paclitaxel must only correspond to Formula I, whatever is the nature of  $R_1$  to  $R_{17}$  substituents.

10 The term "pro-taxanes" used in accordance with the present invention is intended to mean any precursors of any taxanes in the biosynthesis pathway of paclitaxel in plant, fungi and bacteria, including, without limitation, 10-deacetylbaccatin III, baccatin  
15 III, cephalomannine, taxinines, taxuspines, taxusin, 7-xylosyl taxol, 7-epi-10-deacetyl taxol, 10-deacetyl taxol, paclitaxel, 7-epitaxol, taxadiene, geranyl-geranyl-pyrophosphate (GGPP) and farnesyl-pyrophosphate.

20 Exemplary pro-taxanes which may be biotransformed by the bacteria of the present invention include but are not limited to those of the preceding formula I where  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$  are defined in Table 1 below.

25 Exemplary biotransformed taxanes of the present invention include, but are not limited to, those of the


following Formula I where  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ ,  $R_7$ ,  $R_8$ ,  $R_9$ ,  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{14}$ ,  $R_{15}$ ,  $R_{16}$ ,  $R_{17}$  are defined in Table 1.

Table 1


Compound	R <sub>1</sub> <sup>(1)</sup>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub> <sup>(3)</sup>	R <sub>7</sub> <sup>(2)</sup>	R <sub>8</sub> R <sub>17</sub>	R <sub>9</sub> R <sub>10</sub> <sup>(4)</sup>	R <sub>11</sub>	R <sub>12</sub> <sup>(6)</sup>	R <sub>13</sub>	R <sub>14</sub>	R <sub>15</sub> <sup>(5)</sup>	R <sub>16</sub>
1) pacitaxel	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
2) 10-deacetyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
3) 7-epitaxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
4) 10-deacetyl-7- epitaxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
5) 7-epi- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
6) baccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
7) 10-deacetyl baccatin III	α-OH	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
8) cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
9) 10-deacetyl taxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
10) 7-xylosyl taxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
11) 7-xylosyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
12) taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
13) δ-benzoyloxy- taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-benzoyl- oxymethyl	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
14) 9-acetyloxy- taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
15) 9-hydroxy-taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-OH	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
16) taxane Ia	tax	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
17) taxane Ib	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
18) taxane Ic	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-acetyloxy	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
19) taxane Id	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-OH	epoxide	α-acetyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
20) 7-epibaccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>

## Footnotes

(1) "ceph" denotes




"tax" denotes




5

"taxsub" denotes



(2) "xylosyl" denotes




10 (3) " $\alpha$ " denotes the stereoposition of a stereomoiety below the plane of the taxane ring structure shown above




15 " $\beta$ " denotes the stereoposition of a moiety above the plane of the taxane ring structure shown above



(4) "oxetane" denotes the moiety



20 which is




5) "cyclo" denotes the cyclic group formed by bonding the group






to the taxane

A ring as follows:



6) "epoxide" denotes the moiety



5

which is



The term "taxol-associated cytotoxic biological activity" is intended to mean a cytotoxic activity which is sufficient to promote the assembly of tubulin and stabilizes the resulting microtubules of cancer cells causing the division of the cells in two equal daughter cells to be interrupted; and sufficient to cause a disruption in the dynamic equilibrium which exists between microtubules and their depolymerized tubulin dimers, thus preventing completion of the mitotic step which causes a lethal metaphase arrest of cancer cells.

The expression "cancer cells" is intended to mean any cancer cells which include without limitation, ovarian, breast, lung, head and neck cancer cells.

The term "growth supporting nutrient medium" is intended to mean any culture media which include, without limitation, carbon sources, nitrogen sources, amino acids, vitamins and minerals.

25

The term "intercalating agent" is intended to mean any molecule binding to the double stranded DNA structure which include, without limitation, daunorubicine, ethidium bromide, acridine orange, acriflavine  
5 and epirubicine.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1A shows the elution profile, according to HPLC method no. 1, of an organic supernatant extract of  
10 a bacteria in accordance with one embodiment of the present invention, referred to as *Sphingomonas taxi*;

Fig. 1B illustrates the UV spectra of paclitaxel obtained from a standard and of an organic extract of supernatant of a bacteria in accordance with one  
15 embodiment of the present invention, referred to as *Sphingomonas taxi*;

Fig. 2A illustrates a typical chromatogram of an organic extract of the supernatant of a bacteria in accordance with one embodiment of the present invention, referred to as *Bacillus cereus* ssp. *taxi* using  
20 HPLC method no. 2;

Fig. 2B illustrates the ultraviolet spectrum of paclitaxel produced by *Bacillus cereus* ssp. *taxi* compared to a paclitaxel standard;

25 Fig. 2C illustrates the ultraviolet spectrum of 7-xylosyl-10-deacetyltaxol produced by *Bacillus cereus* ssp. *taxi* compared to an authentic standard co-eluting with taxinine M;

Fig. 3A illustrates a typical chromatogram of an  
30 organic extract of the supernatant of a bacteria in accordance with one embodiment of the present invention, referred to as *Pantoea* sp. BCM 1 using HPLC

method no. 2;

Fig. 3B illustrates the ultraviolet spectrum of 7-epi-10-deacetyltaxol produced by *Pantoea* sp. BCM 1 compared to an authentic standard;

5 Fig. 4A illustrates a typical chromatogram of an organic extract of the supernatant of a bacteria in accordance with one embodiment of the present invention, referred to as *Bacillus megaterium* ssp. *taxi* using HPLC method no. 2;

10 Fig. 4B illustrates the ultraviolet spectrum of the bacterial paclitaxel produced by *Bacillus megaterium* ssp. *taxi* compared to an authentic standard;

Fig. 5A illustrates a typical chromatogram of an organic extract of the supernatant of a bacteria in accordance with one embodiment of the present invention, referred to as *Bacillus cereus* ssp. BCM 4 using HPLC method no. 2;

Fig 5B shows a substance having the characteristic ultraviolet spectrum of a taxane compared to a standard of 7-xylosyl-10-deacetyltaxol co-eluting with taxinine M;

20 Fig. 6 illustrates the cytotoxicity of organic extracts of microorganisms isolated from different species of *Taxus* on CRL-1572 cell line as well as the cytotoxicity of taxanes and paclitaxel negative bacteria (STJ.BRP.K1 and *E. coli* TG1);

Fig. 7A illustrates the mass spectrometry of the substances eluted between 45 and 48 minutes using HPLC method no. 1 from an extract of *Sphingomonas taxi*;

30 Fig. 7B illustrates the mass spectrometry of a paclitaxel standard;

Fig. 8A illustrates the almost complete 16S rRNA

gene sequence of *Sphingomonas taxi* (SEQ ID NO:1);

Fig. 8B illustrates the almost complete 16S rRNA gene sequence of *Bacillus cereus* ssp. *taxi* (SEQ ID NO:2);

5 Fig. 8C illustrates the partial 16S rRNA gene sequence of *Bacillus megaterium* ssp. *taxi* (SEQ ID NO:3);

Fig. 8D illustrates the partial 16S rRNA gene sequence of *Pantoea* sp. BCM 1 (SEQ ID NO:4);

10 Fig. 8E illustrates the partial 16S rRNA gene sequence of *Bacillus cereus* ssp. BCM 4 (SEQ ID NO:5);

Fig. 8F illustrates the partial 16S rRNA gene sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Bacillus subtilis* ssp. *taxi* (SEQ ID NO:6);

Fig. 8G illustrates the partial 16S rRNA gene sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Pantoea* sp. BCM 2 (SEQ ID NO:7);

20 Fig. 8H illustrates the partial 16S rRNA gene sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Pantoea* sp. BCM 3 (SEQ ID NO:8);

Fig. 8I illustrates the partial 16S rRNA gene sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Bacillus megaterium* ssp. BCM 9 (SEQ ID NO:9);

30 Fig. 8J illustrates the partial 16S rRNA gene sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Curtobacterium* sp. BCM 5 (SEQ ID NO:10);

Fig. 8K illustrates the partial 16S rRNA gene

sequence of a bacteria in accordance with one embodiment of the present invention, referred to as *Sphingomonas* sp. BCM 7 (SEQ ID NO:11);

5 Fig. 9A illustrates a typical chromatogram of an organic extract from the supernatant of *Bacillus cereus* ssp. *taxi* producing a specific bacterial taxane using HPLC method no. 2;

10 Fig. 9B illustrates the ultraviolet spectrum of the characteristic taxane produced by *Bacillus cereus* ssp. *taxi*;

Fig. 10A illustrates the GC/MS (EI) spectrum of the purified specific bacterial taxane produced by *Bacillus cereus* ssp. *taxi* in accordance with the present invention;

15 Fig. 10B illustrates the mass-to-charge ratios of ions present in the EI mass spectra of the bacterial taxane produced by *Bacillus cereus* ssp. *taxi*;

20 Fig. 11A shows the evolution in time of two taxanes in the sterile culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis* incubated at 150 rpm at 30°C;

25 Fig. 11B shows the evolution of two taxanes transformed by *Sphingomonas taxi* when cultured in the culture medium S-7 supplemented with 1% of an aqueous extract of *Taxus canadensis*, shaken at 150 rpm at 30°C;

30 Fig. 12 shows the HPLC chromatograms of *S. taxi* cultured in the S-7 culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis* where sample name 8A represents the organic extract of the culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*, sample name S 24 hres represents the organic extract of *Sphingomonas taxi* incu-

bated 24 hours in the culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*, sample name S 48 hres represents the organic extract of *Sphingomonas taxi* incubated 48 hours in the culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*, and sample name S 72 hres represents the organic extract of *Sphingomonas taxi* incubated 72 hours in the culture medium supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*;

10 Fig. 13 compares HPLC chromatograms of the supernatant extracts of *S. taxi* (sample name: S 48 hres), *S. taxi* D200 (sample name: SC 48 hres) and *S. Taxi* D201 (sample name: SCP 48 hres) cultured 48 hours in the medium S-7 supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*, and the organic  
15 extract of the medium S-7 supplemented with 1% v/v of an aqueous extract of *Taxus canadensis* (sample name 8A); and

Figs. 14A-14C show the characteristic ultraviolet  
20 let spectrum of the new biotransformed pro-taxanes.

#### DETAILED DESCRIPTION OF THE INVENTION

Plants are hosts of a variety of microorganisms. The relation between the plant and the microorganism  
25 can be saprophytic, parasitic, endophytic or symbiotic. Whatever the relation, there may be genetic exchange between the species. *Taxus*, such as *Taxus canadensis*, which grows in some regions of the province of Quebec, shows significant amounts of paclitaxel in its needles  
30 and stems. Samples of *Taxus canadensis* from seven (7) regions of the province of Quebec were chosen as well as samples of different species of *Taxus* such as *Taxus*



*brevifolia*, *T. cuspidata*, *T. baccata*, *T. hunnewelliana*. Several different bacteria of different genus, such as *Sphingomonas*, *Bacillus*, *Pantoea*, and *Curtobacterium* were isolated from inner parts of samples from  
5 different species of *Taxus*, and all demonstrated taxanes and/or paclitaxel-producing properties.

Bacteria described above, produced taxanes and paclitaxel in fermentation procedures. Bacteria are cultured in a appropriate growth supporting nutrient  
10 medium containing ingredients known to those skilled in the art for the cultivation of microorganisms. Specific examples of appropriate such media are given below. Temperature of cultivation ranges from 10°C to 35°C, and aerobic cultivation is generally preferred.

15 Taxanes and paclitaxel are generally excreted in the supernatant, up to 10% of those substances remain in the cell. Taxanes and paclitaxel thereof may be extracted by performing an organic extraction with an appropriate solvents such as methylene chloride or  
20 ethyl acetate.

In accordance with the present invention, various bacteria producing taxanes and paclitaxel were isolated from different species of *Taxus*.

One bacterium isolated from *Taxus canadensis*  
25 allows for the production of taxanes and paclitaxel at a yield of 1 µg/L, referred to as *Sphingomonas taxi*, has been already deposited at the American Type Culture Collection but was identified as *Erwinia taxi* (ATCC, 12301 Parklawn Drive, Rockville, MD 20852 USA) on April  
30 25, 1995 under deposit number ATCC 55669. The deposit is also available as required by Foreign Patent laws in

countries wherein counterpart applications are filed.

The other strains of the present invention have been deposited at the American Type Culture Collection (ATCC, 12301 Parklawn Drive, Rockville, MD 20852 USA) on December 18, 1997 under deposit accession numbers as set forth below. The deposit are also available as required by Foreign Patent laws in countries wherein counterpart applications are filed.

Strain	ATCC No.
<i>Bacillus cereus</i> ssp. <i>taxi</i>	202061
<i>Bacillus megaterium</i> ssp. <i>taxi</i>	202062
<i>Curtobacterium</i> sp. BCM5	202063
<i>Pantoea</i> sp. BCM2	202064
<i>Bacillus megaterium</i> BCM9	202065
<i>Bacillus cereus</i> BCM4	202066
<i>Sphingomonas</i> <i>taxi</i> D201	202067
<i>Sphingomonas</i> <i>taxi</i> D200	202068
<i>Sphingomonas</i> sp. BCM7	202069
<i>Pantoea</i> sp. BCM3	202070
<i>Pantoea</i> sp. BCM1	202071
<i>Bacillus subtilis</i> ssp. <i>taxi</i>	202072

10

In accordance with one embodiment of the present invention, the bacteria isolated from different species of *Taxus* allow for the production of taxanes and paclitaxel thereof at a yield of 1 to 25 µg/L.

15

In accordance with the present invention, the bacteria isolated from different species of *Taxus* may be employed for the biotransformation of pro-taxanes.

### Isolation of the different microorganisms producing taxanes and paclitaxel

Each plant was divided into 5 parts; needles, twigs, stems, bark and roots. Each inner part of the  
5 plant was verified for the presence of taxanes and paclitaxel producing microorganisms.

The surface of every part of the plant was sterilized with 95% ethanol and then, cut into small pieces with a sterile blade. Pieces were homogenized in sterile  
10 ile water with a POLYTRON™ that had also been sterilized with ethanol 95%. The resulting mix was used to inoculate two different culture media; R2A agar (Difco) and Sabouraud agar (Difco).

Each plate was incubated at 22°C and examined on  
15 a day-to-day basis. The morphology of each colony was meticulously noted and the bacteria were transferred on different media until a pure culture was obtained. A Gram coloration of every bacteria was done before the culture was frozen at -80°C.

20 Over 50 bacteria were isolated from different samples of *Taxus canadensis* of the province of Quebec. In addition, over 30 different bacteria were isolated from different species of *Taxus* which include, without limitations, *Taxus brevifolia*, *T. baccata*, *T. cuspidata*,  
25 *T. hunnewelliana*. Some of them, showing taxanes and paclitaxel production capacities, will be fully described below.

### Screening of microorganisms

In order to verify the production of taxanes and  
30 paclitaxel by microorganisms, each organism was cultured in at least 500 ml of a growth supporting nutrient medium. Any liquid medium allowing taxanes and

paclitaxel thereof production may be employed. Exemplary liquid media are S-7 media (Table 2), and defined media for *Bacillus* (Table 3). Every culture was performed in culture flasks and incubated at a temperature ranging from 20°C to 35°C with constant shaking until a sufficient growth was achieved, generally 18 to 72 hres.

**Table 2**  
**Composition of S-7 medium**

Compounds	g/L
glucose	1
fructose	3
sucrose	6
sodium acetate	1
soytone	1
thiamine	0,001
biotine	0,001
pyridoxal-HCl	0,001
Ca pantothenate	0,001
MgSO <sub>4</sub>	0,0036
CaNO <sub>3</sub>	0,0065
Cu (NO <sub>3</sub> ) <sub>2</sub>	0,001
ZnSO <sub>4</sub>	0,0025
MnCl <sub>2</sub>	0,005
FeCl <sub>3</sub>	0,002
phenylalanine	0,005
sodium benzoate	0,1
KH <sub>2</sub> PO <sub>4</sub> 1M (pH 6,8)	1 ml

Table 3Composition of the defined medium for *Bacillus*

Compounds	g/L
L-glutamic acid	10
glucose	5
citric acid	1
K <sub>2</sub> HPO <sub>4</sub>	0,5
KH <sub>2</sub> PO <sub>4</sub>	0,5
MgSO <sub>4</sub> -7H <sub>2</sub> O	0,2
MnSO <sub>4</sub> -4H <sub>2</sub> O	0,01
FeSO <sub>4</sub> -7H <sub>2</sub> O	0,01

The culture was then centrifuged and the pellet  
5 separated from the supernatant by decantation. To ver-  
ify if taxanes and paclitaxel were secreted in the  
medium or if it was confined within the cells, both  
were tested for the presence of the drug. Since tax-  
anes and paclitaxel are hydrophobic, and in order to  
10 concentrate each sample, an extraction with an organic  
solvent was performed. For the pellet, the cells were  
dried and about 200 mg were powdered and ultrasonicated  
twice for 40 minutes in 3 ml methanol. The extracts  
were dried at 25°C. The residue was dissolved by add-  
15 ing 2 ml of methylene chloride and 2ml of distilled  
water. After appropriate shaking, the mixture was cen-  
trifuged at 4 000 rpm for 5 min. The methylene chlo-  
ride fraction was collected and dried under reduced  
pressure. Finally, the residue is dissolved in 0,5 ml  
20 of HPLC grade methanol.

The supernatant is extracted with one volume of  
methylene chloride. After appropriate shaking, the

organic fraction is evaporated to dryness under reduced pressure. The residue is then resolubilized in 50 ml of methylene chloride and 50 ml of distilled water. After appropriate shaking, each fraction was collected  
5 and dried under reduced pressure. Each residue is dissolved in a measured minimal volume of HPLC grade methanol. All samples were kept frozen at -20°C.

**a) HPLC screening**

10 HPLC method no. 1

Some extracts were analyzed by High Performance Liquid Chromatography (HPLC) on a system consisting of a WATERS™ 625 LC pump, a WATERS™ 996 photodiode array spectrophotometer, and a WATERS™ 717plus autosampler.  
15 Chromatography was performed with a phenyl column from Waters ( 5µm particle size, 6 mm X 15 mm) with a guard module. The injection volume varies from 50 to 150 µl and the flow rate maintained at 1 ml/min. The following elution program was used;  
20 0 to 20 min.: methanol:water:acetonitrile  
(20:65:15) ramped to methanol:water:acetonitrile  
(20:45:35)  
20 to 50 min.: methanol:water:acetonitrile (20:45:35)  
ramped to methanol:water:acetonitrile  
25 (20:25:55)  
50 to 60 min.: methanol:water:acetonitrile (20:25:55)  
ramped to methanol 100%

Table 4 identifies the retention times of known authentic standards on HPLC methods no. 1 and no. 2.  
30 Using HPLC no. 1, paclitaxel has a retention time of 46 minutes. In Fig. 1, we show the ultraviolet spectrum



of paclitaxel produced by *Sphingomonas taxi*. The spectrum is very characteristic with a second maximum of absorption at 230nm. This figure illustrates that *Sphingomonas taxi* produces a compound having the same retention time and the same UV spectrum as paclitaxel.

HPLC method no. 2

Some extracts were analyzed on the same HPLC system with a curosil-PFP column (250 mm X 3.2 mm) from Phenomenex with a guard module. Injections varied from 50 ul to 150 ul and the flow rate maintained at 0.8 ml/min. The following gradient program was used;

0 to 50 min.: acetonitrile:water (25:75) ramped to  
acetonitrile:water (65:35)

50 to 62.5 min.: acetonitrile:water (65:35) ramped to  
methanol 100%

62.5 to 65 min.: methanol 100% to acetonitrile:water  
(25:75)

65 to 75 min.: acetonitrile:water (25:75)

As shown in Table 4, using HPLC method no. 2, paclitaxel is eluted at 36.987 minutes.

Table 4

Retention time of taxanes standards using HPLC methods  
no. 1 and no. 2

Taxanes	Retention time using	
	HPLC method no. 1	HPLC method no. 2
10-deacetyl baccatin III	n/a	12.037 min.
baccatin III	n/a	20.670 min.
7-xylosyl-10-deacetyltaxol B	n/a	24.870 min.
7-xylosyl-10-deacetyltaxol and taxinine M	n/a	27.120 min.
7-xylosyl-10-deacetyltaxol C	n/a	28.770 min.
10-deacetyltaxol and 7-xylosyltaxol	n/a	30.770 min.
cephalomannine	n/a	34.753 min.
7-epi-10-deacetyltaxol	n/a	35.703 min.
paclitaxel	46 minutes	36.987 min.
taxol C	n/a	38.853 min.
7-epitaxol	n/a	42.287 min.

5            Fig. 2A shows a typical chromatogram of an organic extract of *Bacillus cereus* ssp. *taxi*, and in Figs. 2B and 2C there is compared the UV spectra of the substances produced by *Bacillus* with authentic commercial plant standards. This figure clearly illustrates  
10 the ability of *Bacillus cereus* ssp. *taxi* to produce paclitaxel and 7-xylosyl-10-deacetyltaxol.

Fig. 3A shows the typical HPLC chromatogram of the supernatant of *Pantoea* sp. BCM 1 and, in Fig. 3B there is compared the bacterial 7-epi-10-deacetyltaxol  
15 against an authentic commercial plant standard, establishing the production of 7-epi-10-deacetyltaxol by *Pantoea* sp. BCM 1.

Fig. 4A shows a typical chromatogram of *Bacillus megaterium* ssp. *taxi* and, in Fig. 4B there is compared  
20 the ultraviolet spectrum of the bacterial paclitaxel

with an authentic standard proving the capacity of this bacterium to produce paclitaxel.

Fig. 5A shows a typical chromatogram of *Bacillus cereus* ssp. BCM 4 and, Fig 5B shows a substance having the characteristic ultraviolet spectrum of a taxane compared to a standard of 7-xylosyl-10-deacetyltaxol co-eluting with taxinine M, proving the capacity of *Bacillus cereus* ssp. BCM 4 to produce taxanes.

**b) Cytotoxicity on cancer cells**

10 An ovarian cancer cell line obtained from American Type Culture Collection (12301 Parklawn Drive, Rockville, MD 20852 USA) under ATCC accession number CRL-1572) was chosen for the investigation. Briefly, 2 000 cells/well of a 96-well microplate were inoculated. 15 After 2 days, different dilutions of the drug were added in a volume of 100 µl. Three days later, the viability of cells was measured with ALAMAR™ blue (B. Pagé et al. 1993, *Intl. J. of Oncology*, 3:473-476). The ATCC CRL-1572 cell line is particularly sensitive 20 to paclitaxel (ID<sub>50</sub> of 1 ng/ml). Microbial extracts have also been tested for their cytotoxicity on those cells. Fig. 6 shows the cytotoxicity of different bacterial supernatant extracts. This figure clearly demonstrates that culture supernatant extracts from *Sphingomonas taxi* and *Bacillus cereus* ssp. *taxi* are at least 25 5-fold more cytotoxic than extracts from non-paclitaxel producing bacteria such as STJ.BRP.K1 and *E. coli* TG1.

**c) Mass spectrometry**

30 Mass spectrometry of the substances eluted between 45 and 48 minutes with HPLC method no.1 from an

extract of *Sphingomonas taxi* was performed. Fig. 7A shows the results obtained. In every sample, a substance with a molecular weight (M.W.) of 853,5 daltons appears. The theoretical M.W. of paclitaxel is 854 daltons. Even if these extracts were partially purified, essentially the major compound obtained is paclitaxel.

**d) Characterization of a bacterial taxane by EI/MS**

Fig. 9A, shows a typical chromatogram of *Bacillus cereus* ssp. *taxi* producing a characteristic taxane having a retention time of 15 minutes using HPLC method no. 2. In Fig. 9B, the characteristic ultraviolet spectrum of this substance is illustrated. It was purified on HPLC using HPLC method no. 2 and, analyzed in Electro Ionization Mass Spectrometry (EI/MS) as described by Thomas D. McClure et al. (*J. Am. Soc. Mass Spectrom.*, 1992, 3, pp. 672-679).

Fig. 10A shows the EI spectrum of this taxane and Fig. 10B shows the characteristic fragments of this taxane. Since this taxane has never been observed in plant extracts of different species of *Taxus*, and this compound is produced by at least two species of *Bacillus* (*cereus* ssp. *taxi* and *megaterium* ssp. *taxi*), we consider this novel taxane unique to microorganisms isolated from *Taxus*.

**Characterization of taxane and paclitaxel producing microorganisms**

**a) Biochemical and physiological characteristics**

Table 5 shows the species of *Taxus* from which each bacterium was isolated, the Gram coloration, the

morphology of colonies on different culture media and, some biochemical characteristics. This Table clearly illustrates the diversity of our taxanes and paclitaxel producing bacteria i.e. we have isolated and identified

5 Gram-negative and Gram-positive rods including Actinomyces from different species of *Taxus* which all possess taxanes and/or paclitaxel producing capacities.

**Table 5**

**Cellular, morphological and biochemical characteristics  
of some taxanes and paclitaxel producing bacteria**

Name	Isolated from	Gram	Cellular morphology	Description of colonies on blood agar	Description of colonies on TA-1	Catalase	Urease
<i>Sphingomonas taxi</i>	<i>Taxus canadensis</i>	-	Rod	No growth	3 days to grow Orange-Opaque Glossy-Convex Circular contour	+	-
<i>Bacillus cereus ssp. taxi</i>	<i>Taxus canadensis</i>	+	Rod	Double hemolysis Gray-green- Opaque Dry-Dull-Flat Irregular edge	Cream- Opaque Dry-Dull-Flat Irregular edge	+	-
<i>Bacillus megaterium ssp. taxi</i>	<i>Taxus hunnewelliana</i>	+	Rod	Gray- Opaque Dry-Dull Convex Irregular edge	Cream- Opaque Glossy Slightly convex Irregular edge	+	-
<i>Pantoea sp. BCM 1</i>	<i>Taxus hunnewelliana</i>	-	Rod	Semi-translucent Glossy-Flat Regular edge	Cream with yellow pigment in middle Opaque-Glossy- Flat Regular edge	+	-
<i>Pantoea sp. BCM 2</i>	<i>Taxus cuspidata</i>	-	Rod	Yellow Opaque Glossy-Flat Regular edge	Yellow Semi-translucent Glossy-Flat Regular edge	+	-
<i>Pantoea sp. BCM 3</i>	<i>Taxus cuspidata</i>	-	Rod	Semi-translucent Glossy-Flat Regular edge	Yellow Semi-translucent Glossy-Flat Regular edge	+	-
<i>Bacillus cereus ssp. BCM 4</i>	<i>Taxus brevifolia</i>	+	Rod	Double hemolysis Gray-green- Opaque Dry-Dull-Flat Irregular edge	Cream- Opaque Dry-Dull-Flat Irregular edge	+	-
<i>Bacillus subtilis ssp. taxi</i>	<i>Taxus baccata</i>	+	Rod	Semi-translucent Dry, Dull Convex Regular edge	Cream- Translucent Dry, Dull Convex Regular edge	+	-
<i>Bacillus megaterium ssp. BCM 9</i>	<i>Taxus hunnewelliana</i>	+	Rod	Gray- Opaque Dry-Dull Convex Irregular edge	Cream- Opaque Glossy Slightly convex Irregular edge	+	-
<i>Curtobacterium sp. BCM 5</i>	<i>Taxus brevifolia</i>	+	Rod	2 days to grow Cream Semi-translucent Glossy-Flat Regular edge	2 days to grow Yellow Semi-translucent Glossy-Flat Regular edge	+	-
<i>Sphingomonas sp. BCM 7</i>	<i>Taxus hunnewelliana</i>	-	Rod	No growth	3 days to grow Orange-Opaque Glossy-Convex Regular edge	+	-

5

**b) Identification of the genus of taxanes and paclitaxel producing microorganisms**

The genus of each taxanes and paclitaxel producing bacteria was determined by sequencing the 16S



rRNA genes. Genomic DNA of each strain was used as template for PCR (Polymerase Chain Reaction). Primers based on conserved regions at the beginning and the end of the 16S rRNA gene, SSU-27 (5' -  
5 AGAGTTTGATCMTGGCTCAG-3'; SEQ ID NO:12), and SSU-1 492 (5' -TACGGYTACCTTGTTACGACTT-3'; SEQ ID NO:13), were used to amplify a portion of the 16S gene. The amplicons were purified with the "PCR purification kit" (sold by Qiagen) and sequenced using the ABI Prism System.  
10 Sequence analysis was performed using GCG software package (Genetics Computer Group Inc., Madison, WI).

Fig. 8A shows the almost complete sequence of the 16S rRNA gene of *Sphingomonas taxi*. Since this strain has unique biosynthetic capacities and more than  
15 3% sequence difference with the 16S rRNA genes of other known species of *Sphingomonas*, we created a new species and named it *taxi* on the behalf of its isolation source. Fig. 8B shows the almost complete sequence of the 16S rRNA gene of *Bacillus cereus* ssp. *taxi*. Since  
20 this bacterium possesses unique metabolic capacities, and in order to differentiate this species from other known *Bacillus cereus*, we identified it by subspecies name *taxi* also on the behalf of its isolation source. In Figs. 8C to 8L, we show partial sequences of the 16S  
25 rRNA genes of other taxanes and/or paclitaxel producing microorganisms.

Consequently, in accordance with the present invention, a plurality of bacteria isolated from different species of *Taxus* can be used for the mass pro-  
30 duction of paclitaxel and other taxanes. Based on the analysis of partial 16S rRNA gene sequences, and morphological and biochemical characteristics, we assigned

the following genera, species, and subspecies or strain names to our paclitaxel and taxanes producing bacterial isolates; *Sphingomonas taxi*, *Bacillus cereus* ssp. *taxi*, *Bacillus megaterium* ssp. *taxi*, *Pantoea* sp. BCM 1, *Pantoea* sp. BCM 2, *Pantoea* sp. BCM 3, *Bacillus cereus* ssp. BCM 4, *Bacillus subtilis* ssp. *taxi*, *Bacillus megaterium* ssp. BCM 9, *Curtobacterium* sp. BCM 5 and *Sphingomonas* sp. BCM 7.

10 **Biotransformation of taxanes**

a) **Preparation of an aqueous extract of *Taxus canadensis***

Fresh cuttings of needles and small twigs (10 g) of a sample of *Taxus canadensis* are homogenized in 100 ml of distilled water. The solution is then centrifuged at 7 000 rpm and the clear supernatant sterilized by filtration on a 0,22  $\mu$ m filter. The solution is kept frozen at -20°C until utilization.

20 b) **Biotransformation of taxanes by taxanes and paclitaxel producing bacteria**

The growth supporting nutrient medium S-7 is supplemented with 1% v/v on an aqueous extract of *Taxus canadensis*. This resulting supplemented medium is then inoculated with a thawed vial of a pure culture of one of our strain and incubated at 30°C with constant shaking for a time sufficient to allow biotransformation of pro-taxanes.

The culture is then centrifuged and the remaining supernatant extracted with one volume of methylene chloride or ethyl acetate. After appropriate shaking, the organic fraction is evaporated to dryness under reduced pressure. The residue is then resolubilized in

50 ml of methylene chloride or ethyl acetate, and 50 ml of distilled water. After appropriate shaking, each fraction was collected and dried under reduced pressure. Each residues is dissolved in a measured minimal  
5 volume of HPLC grade methanol. All samples were kept frozen at -20°C until analysis.

Samples were analyzed on HPLC using method no. 2. Fig. 11 shows the evolution of two taxanes in A) the sterile culture medium S-7 supplemented with 1% v/v  
10 of an aqueous extract of *Taxus canadensis* shaken at 150 rpm and incubated at 30°C, and in B) the supernatant of *Sphingomonas taxi* cultured in the culture medium S-7 supplemented with 1% v/v of an aqueous extract of *Taxus canadensis* shaken at 150 rpm and incubated at 30°C.  
15 This figure clearly illustrates that in the supernatant of *Sphingomonas taxi*, the diminution of the taxane eluted at 12 minutes corresponds to the proportional elevation of the taxane eluted at 24 minutes, proving the capacity of *Sphingomonas taxi* to biotransform tax-  
20 anes.

In addition, Fig. 12 compares the HPLC chromatogram of the organic extract of the culture supernatant of *S. taxi* incubated 24, 48 and 72 hours in the culture medium S-7 supplemented with 1% v/v of an aque-  
25 ous extract of *Taxus canadensis*. This Fig. 12 clearly illustrates the production of pro-taxanes. The ultra-violet spectrum of one of these pro-taxanes is illustrated in Fig. 14A. Fig. 14A compares the UV spectrum of the new biotransformed taxane produced by *S. taxi*  
30 with taxanes from the aqueous extract of *Taxus canadensis*.

**Mutagenesis of taxanes and paclitaxel producing bacteria**

Typically, 20 ml of the culture medium TA-1 with 200 µg/ml of daunorubicin (purchased from Rhône-Poulenc) were inoculated with 500 µl of an overnight culture. The resulting broth was incubated at 200 rpm at 30°C for 2 days. After this time 10 ml of the broth are added to 10 ml of fresh medium containing 200 µg/ml of daunorubicin and incubated as described above. The preceding step is repeated as necessary to obtain mutated bacteria. Those mutants were further isolated on the solid culture medium TA-1 (composition as follows).

**Solid culture medium TA-1**

Ingredient	amount
glucose	5 g
tryptone	20 g
yeast extract	5 g
NaCl	0.5 g
agar	15 g
H <sub>2</sub> O	1 L

15

**Biotransformation of taxanes by mutated strains**

The growth supporting nutrient medium S-7 is supplemented with 1% v/v on an aqueous extract of *Taxus canadensis*. This resulting supplemented medium is then inoculated with a thawed vial of a pure culture of one of our strain and incubated at 30°C with constant shaking for a time sufficient to allow biotransformation of pro-taxanes.

The culture is then centrifuged and the remaining supernatant extracted with one volume of methylene

25

chloride or ethyl acetate. After appropriate shaking, the organic fraction is evaporated to dryness under reduced pressure. The residue is then resolubilized in 50 ml of methylene chloride or ethyl acetate, and 50 ml of distilled water. After appropriate shaking, each fraction was collected and dried under reduced pressure. Each residue is dissolved in a measured minimal volume of HPLC grade methanol. All samples were kept frozen at -20°C until analysis.

10           Samples were analyzed on HPLC using method no. 2. Fig. 13 shows HPLC chromatograms of *S. taxi*, *S. taxi* D200 and *S. taxi* D201 incubated the same time (48 hours) in the culture medium S-7 supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*. All  
15           cultures had comparable cell density. This figure clearly illustrates the improved yields of biotransformation by the mutated strains *S. taxi* D200 and *S. taxi* D201. In Figs. 14B and 14C the characteristic ultra-violet spectrum of the new pro-taxanes, produced by the  
20           mutated strains *S. taxi* D200 and *S. taxi* D201, are compared with the UV spectrum of two taxanes from *Taxus canadensis*.

          The present invention will be more readily understood by referring to the following examples which  
25           are given to illustrate the invention rather to limit its scope.

#### EXAMPLE I

##### **Mass production of paclitaxel using *Sphingomonas taxi***

          A colony of a pure culture of *Sphingomonas taxi*  
30           is used to inoculate 5 ml of S-7 culture medium. The broth is incubated 2-3 days with constant shaking (90 rpm) at 22°C. This 5 ml is then transferred into 5

liters of the same culture medium. The resulting broth is incubated as described above in aerobic conditions.

After 4-5 days of incubation, or after the maximum cell density is reached, the cell pellet is separated by centrifugation. Hydrophobic compounds are then extracted from the supernatant by partition with one volume of dichloromethane. Each organic fraction is evaporated to dryness and, the residue is solubilized in a minimal amount of HPLC grade methanol, typically 500  $\mu$ l to 1 ml.

Paclitaxel and taxanes are further purified by HPLC on a phenyl column using HPLC method no. 1. Typically, up to 400  $\mu$ l of the methanolic solution are injected and fractions of 0,5 ml to 1 ml are collected. Fractions containing paclitaxel or taxanes are evaporated to dryness.

Using this method, from 200 ng to 1  $\mu$ g of paclitaxel per liter of culture medium were purified.

20

#### EXAMPLE II

##### **Mass production of taxanes and paclitaxel using *Bacillus cereus* ssp. *taxi***

A thawed vial of a pure dense cell suspension of *Bacillus cereus* ssp. *taxi* is used to inoculate 500 ml of the defined medium for *Bacillus*. The broth is incubated 1 to 3 days with constant shaking (150 rpm) at 30°C. The cell pellet is then separated from the supernatant by centrifugation. Hydrophobic substances are extracted from the supernatant by an extraction with one volume of dichloromethane. The organic fraction is evaporated to dryness under reduced pressure,

and residues resolubilized in a minimal amount of HPLC grade methanol.

Taxanes and paclitaxel are further purified on HPLC using method no. 2. Using this method, from 0,2 to 10 µg of paclitaxel can be produced, and from 0,2 to 15 µg of taxanes, including the specific bacterial taxane illustrated in Fig. 13B, can also be produced.

#### EXAMPLE III

##### 10 Mass production of taxanes and paclitaxel using *Bacillus megaterium* ssp. *taxi*

A thawed vial of a pure dense cell suspension of *Bacillus megaterium* ssp. *taxi* is used to inoculate 500 ml of the S-7 culture medium. The broth is incubated 1 to 3 days with constant shaking (150 rpm) at 30°C. The cell pellet is then separated from the supernatant by centrifugation. Hydrophobic substances are extracted from the supernatant by an extraction with one volume of dichloromethane. The organic fraction is evaporated to dryness under reduced pressure, and residues resolubilized in a minimal amount of HPLC grade methanol.

Taxanes and paclitaxel are further purified on HPLC using method no. 2. Using this method, from 1 to 12 µg of paclitaxel can be produced, and from 1 to 15 µg of taxanes, including the specific bacterial taxane illustrated in Fig. 13B, can also be produced.

#### EXAMPLE IV

##### 30 Biotransformation of pro-taxanes by *Sphingomonas taxi*

The growth supporting nutrient medium S-7 is supplemented with 1% v/v on an aqueous extract of *Taxus*



*canadensis*. This resulting supplemented medium is then inoculated with a thawed vial of a pure culture of one of *Sphingomonas taxi* and incubated at 30°C with constant shaking for 24 to 96 hours.

5           The culture is then centrifuged and the remaining supernatant extracted with one volume of methylene chloride. After appropriate shaking, the organic fraction is evaporated to dryness under reduced pressure. The residue is then resolubilized in 50 ml of methylene  
10 chloride and 50 ml of distilled water. After appropriate shaking, the organic fraction is collected and dried under reduced pressure. The residue is dissolved in 500µl of HPLC grade methanol and 100 µl of the methanolic solution are analyzed on HPLC using method  
15 no. 2 and compared to the resulting chromatogram of the organic extract of the growth-supporting nutrient medium supplemented with and aqueous extract of *Taxus canadensis* 1% v/v shaken the same time.

          As illustrated in Figs. 11, 12 and 14A, *Sphingomonas taxi* is able to biotransform taxanes into new  
20 pro-taxanes.

#### EXAMPLE V

##### **Mutagenesis of *Sphingomonas taxi***

25           20 ml of the culture medium TA-1 with 200 µg/ml of daunorubicin (purchased from Rhône-Poulenc) were inoculated with 500 µl of an overnight culture of *Sphingomonas taxi*. The resulting broth was incubated at 200 rpm at 30°C for 2 days. After this time 10 ml  
30 of the broth are added to 10 ml of fresh medium containing 200 µg/ml of daunorubicin and incubated as described above. The preceding step is repeated as necessary to obtain mutated bacteria. Those mutants were further isolated on the solid culture medium TA-1. Two

new mutated strains were obtained named *Sphingomonas taxi* D200 and *Sphingomonas taxi* D201.

#### EXAMPLE VI

##### 5 **Biotransformation of taxanes by *Sphingomonas taxi* D200 and *Sphingomonas taxi* D201**

The growth supporting nutrient medium S-7 is supplemented with 1% v/v on an aqueous extract of *Taxus canadensis*. This resulting supplemented medium is then  
10 inoculated with a thawed vial of a pure culture of one of our mutated strain and incubated at 30°C at 150 rpm. Cultures were stopped after 24, 48 and 72 hours of incubation.

Cultures were then centrifuged and the remaining  
15 supernatants extracted with one volume of ethyl acetate. After appropriate shaking, the organic fractions were evaporated to dryness under reduced pressure. Residues were then resolubilized in 50 ml of ethyl acetate. After appropriate shaking, fractions  
20 were collected and dried under reduced pressure. Each residues were dissolved in a measured minimal volume of HPLC grade methanol. All samples were kept frozen at -20°C until analysis.

Samples were analyzed on HPLC using method no.  
25 2. Fig. 13 shows HPLC chromatograms of *S. taxi*, *S. taxi* D200 and *S. taxi* D201 incubated the same time (48 hours) in the culture medium S-7 supplemented with 1% v/v of an aqueous extract of *Taxus canadensis*. All cultures had comparable cell densities. This figure  
30 clearly illustrates the improved yields of biotransformation by the mutated strains *S. taxi* D200 and *S. taxi* D201. In Fig. 14B the characteristic ultraviolet spec-

trum of the new pro-taxanes produced by *S. taxi* D200 is compared with the UV spectrum of two taxanes from *Taxus canadensis*. Fig. 14C compares the UV spectrum of the new biotransformed taxane produce by *S. taxi* D201 with  
5 taxanes from the aqueous extract of *Taxus canadensis*.

While the invention has been described in connection with specific embodiments thereof, it will be understood that is capable of further modifications and this application is intended to cover any variations,  
10 uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as many be applied to  
15 the essential features hereinbefore set forth, and as follows in the scope of the appended claims.



# American Type Culture Collection

12301 Parklawn Drive • Rockville, MD 20852 USA • Telephone: 301-231-5519 or 231-5532 • FAX: 301-816-4366

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF  
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

## INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3  
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2

To: (Name and Address of Depositor or Attorney)

BCM Developpement Inc.  
Attn: Nathalie Landry  
125 rue Dalhousie, suite 218  
Quebec Canada G1K 4C5

REC'D 30 MAR 1999  
WIPO PCT

Deposited on Behalf of: BCM Developpement

Identification Reference by Depositor:

ATCC Designation

<i>Bacillus cereus</i> ssp. taxi	202061
<i>Bacillus megaterium</i> ssp. taxi	202062
<i>Curtobacterium</i> sp. BCM5	202063
<i>Pantoea</i> sp. BCM2	202064
<i>Bacillus megaterium</i> BCM9	202065
<i>Bacillus cereus</i> BCM4	202066

The deposits were accompanied by: ☐ a scientific description ☒ a proposed taxonomic description indicated above. The deposits were received December 12, 1997 by this International Depository Authority and have been accepted.

AT YOUR REQUEST: ☒ We will not inform you of requests for the strains.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested December 18, 1997. On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Rockville, Md. 20852 USA

Signature of person having authority to represent ATCC:

Barbara M. Hailey  
Barbara M. Hailey, Administrator, Patent Depository

Date: December 18, 1997

cc: France Cote



# American Type Culture Collection

12301 Parklawn Drive • Rockville, MD 20852 USA • Telephone: 301-231-5519 or 231-5532 • FAX: 301-816-4366

BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF  
THE DEPOSIT OF MICROORGANISMS FOR THE PURPOSES OF PATENT PROCEDURE

## INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT ISSUED PURSUANT TO RULE 7.3  
AND VIABILITY STATEMENT ISSUED PURSUANT TO RULE 10.2

To: (Name and Address of Depositor or Attorney)

BCM Developpement Inc.  
Attn: Nathalie Landry  
125 rue Dalhousie, suite 218  
Quebec Canada G1K 4C5

Deposited on Behalf of: BCM Developpement

Identification Reference by Depositor:

ATCC Designation

<i>Sphingomonas</i> taxi D201	202067
<i>Sphingomonas</i> taxi D200	202068
<i>Sphingomonas</i> sp. BCM7	202069
<i>Pantoea</i> sp. BCM3	202070
<i>Pantoea</i> sp. BCM1	202071
<i>Bacillus subtilis</i> ssp. taxi	202072

The deposits were accompanied by: \_\_\_\_ a scientific description X a proposed taxonomic description indicated above. The deposits were received December 12, 1997 by this International Depository Authority and have been accepted.

AT YOUR REQUEST: X We will not inform you of requests for the strains.

The strains will be made available if a patent office signatory to the Budapest Treaty certifies one's right to receive, or if a U.S. Patent is issued citing the strains, and ATCC is instructed by the United States Patent & Trademark Office or the depositor to release said strains.

If the cultures should die or be destroyed during the effective term of the deposit, it shall be your responsibility to replace them with living cultures of the same.

The strains will be maintained for a period of at least 30 years from date of deposit, or five years after the most recent request for a sample, whichever is longer. The United States and many other countries are signatory to the Budapest Treaty.

The viability of the cultures cited above was tested December 18, 1997. On that date, the cultures were viable.

International Depository Authority: American Type Culture Collection, Rockville, Md. 20852 USA

Signature of person having authority to represent ATCC:

Barbara M. Hailey  
Barbara M. Hailey, Administrator, Patent Depository


Date: December 18, 1997

cc: France Cote

WHAT IS CLAIMED IS:

1. A process for producing a taxane, which comprises the steps of:
  - a) culturing at least one bacteria isolated from plant species of *Taxus* in growth-supporting nutrient medium capable of promoting growth and reproduction of said bacteria, and wherein said culturing is effected for a time sufficient to allow production of a taxane; and
  - b) recovering a taxane from said bacteria or medium of step a).
2. The process of claim 1, wherein the plant species of *Taxus* is of the species selected from the group consisting of but not limited to *Taxus canadensis*, *T. brevifolia*, *T. baccata*, *T. hunnewelliana* or *T. cuspidata*.
3. The process of claim 1, wherein said bacteria is of the genus selected from the group consisting of *Sphingomonas*, *Bacillus*, *Pantoea*, and *Curtobacterium*.
4. The process of claim 1, wherein said taxane produced is paclitaxel.
5. The process of claim 1, wherein said taxane is selected from the group consisting of paclitaxel, 10-deacetylcephalomannine, 7-epitaxol, 10-deacetyl-7-epitaxol, 7-epicephalomannine, baccatin III, 10-deacetyl-baccatin III, cephalomannine, 7-epibaccatin III, 7-

xylosyltaxol, 7-xylosyl-cephalomannine, taxagifine,  $\delta$ -benzoyloxy taxagifine, 9-acetyloxy taxusin, 9-hydroxy taxusin, taxane Ia, taxane Ib, taxane Ic, taxane Id and any taxane corresponding to Formula I:



wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub> are defined in Table 1.



Table 1

Compound	R <sub>1</sub> <sup>(1)</sup>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub> <sup>(3)</sup>	R <sub>7</sub> <sup>(2)</sup>	R <sub>8</sub> R <sub>17</sub>	R <sub>9</sub> R <sub>10</sub> <sup>(4)</sup>	R <sub>11</sub>	R <sub>12</sub> <sup>(6)</sup>	R <sub>13</sub>	R <sub>14</sub>	R <sub>15</sub> <sup>(5)</sup>	R <sub>16</sub>
1) paclitaxel	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
2) 10-deacetyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
3) 7-epitaxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
4) 10-deacetyl-7- epitaxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
5) 7-epi- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
6) baccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
7) 10-deacetyl baccatin III	α-OH	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
8) cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
9) 10-deacetyl taxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
10) 7-xylosyl taxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
11) 7-xylosyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
12) taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
13) δ-benzoyloxy- taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-benzoyl- oxymethyl	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
14) 9-acetyloxy- taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
15) 9-hydroxy-taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-OH	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
16) taxane Ia	tax	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
17) taxane Ib	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
18) taxane Ic	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-acetyloxy	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
19) taxane Id	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-OH	epoxide	α-acetyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
20) 7-epibaccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>

SUBSTITUTE SHEET (RULE 26)

6. A bacterial taxane having the characterizing ultraviolet spectrum and the specific retention time shown in Fig. 13, and the EI spectrum shown in Fig. 14A.

7. The process of claim 1, wherein the taxane produced having the characterizing ultraviolet spectrum and the specific retention time shown in Fig. 13, and the EI spectrum shown in Fig. 14A.

8. The process of claim 1, wherein the bacteria is *Bacillus cereus* ssp. *taxi*.

9. The process of claim 1, wherein the bacteria is *Bacillus megaterium* ssp. *taxi*.

10. The process of claim 1, wherein the bacteria is *Pantoea* sp. BCM 1.

11. The process of claim 1, wherein the bacteria is *Pantoea* sp. BCM 2.

12. The process of claim 1, wherein the bacteria is *Pantoea* sp. BCM 3.

13. The process of claim 1, wherein the bacteria is *Bacillus cereus* ssp. BCM 4.

14. The process of claim 1, wherein the bacteria is *Bacillus subtilis* ssp. *taxi*.

15. The process of claim 1, wherein the bacteria is *Bacillus megaterium* ssp. BCM 9.


16. The process of claim 1, wherein the bacteria is *Curtobacterium* sp. BCM 5.

17. The process of claim 1, wherein said growth-supporting nutrient medium include carbon sources, nitrogen sources, amino acids, vitamins and minerals.

18. A process for producing biotransformed taxanes, which comprises the steps of:

- a) culturing at least one bacteria isolated from plant species of *Taxus* in growth-supporting nutrient medium supplemented with pro-taxanes, and wherein said culturing is effected for a time sufficient to allow the biotransformation of pro-taxanes into taxanes; and
- b) recovering at least one from said bacteria or medium of step a).

19. The process of claim 18, wherein said taxane is selected from the group consisting of paclitaxel, 10-deacetylcephalomannine, 7-epitaxol, 10-deacetyl-7-epitaxol, 7-epicephalomannine, baccatin III, 10-deacetyl-baccatin III, cephalomannine, 7-epibaccatin III, 7-xylosyltaxol, 7-xylosyl-cephalomannine, taxagifine,  $\delta$ -benzoyloxy taxagifine, 9-acetyloxy taxusin, 9-hydroxy taxusin, taxane Ia, taxane Ib, taxane Ic, taxane Id and any taxane corresponding to Formula I :



I

wherein R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub>, R<sub>8</sub>, R<sub>9</sub>, R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, R<sub>16</sub>, R<sub>17</sub> are defined in Table 1,

Table 1

Compound	R <sub>1</sub> <sup>(1)</sup>	R <sub>2</sub>	R <sub>3</sub>	R <sub>4</sub>	R <sub>5</sub>	R <sub>6</sub> <sup>(3)</sup>	R <sub>7</sub> <sup>(2)</sup>	R <sub>8</sub> R <sub>17</sub>	R <sub>9</sub> R <sub>10</sub> <sup>(4)</sup>	R <sub>11</sub>	R <sub>12</sub> <sup>(6)</sup>	R <sub>13</sub>	R <sub>14</sub>	R <sub>15</sub> <sup>(5)</sup>	R <sub>16</sub>
1) pacitaxel	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
2) 10-deacetyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
3) 7-epitaxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
4) 10-deacetyl-7- epitaxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
5) 7-epi- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
6) baccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
7) 10-deacetyl baccatin III	α-OH	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
8) cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
9) 10-deacetyl taxol	tax	CH <sub>3</sub>	H	β-OH	=O	β-CH <sub>3</sub>	β-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
10) 7-xylosyl taxol	tax	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
11) 7-xylosyl- cephalomannine	ceph	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	β-xylosyl	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
12) taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
13) 8-benzoyloxy- taxagifine	=O	α-CH <sub>3</sub>	β-OH	β-acetyloxy	α-acetyloxy	β-benzoyl- oxymethyl	β-acetyloxy	H	α-cinnamoyloxy	methylene (=CH <sub>2</sub> )	α-acetyloxy	β-H	H	cyclo	α-CH <sub>3</sub>
14) 9-acetyloxy- taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
15) 9-hydroxy-taxusin	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-OH	β-CH <sub>3</sub>	H	H	α-acetyloxy	methylene (=CH <sub>2</sub> )	H	H	H	CH <sub>3</sub>	CH <sub>3</sub>
16) taxane Ia	tax	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
17) taxane Ib	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
18) taxane Ic	taxsub	CH <sub>3</sub>	H	=O	=O	β-CH <sub>3</sub>	α-acetyloxy	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
19) taxane Id	α-acetyloxy	CH <sub>3</sub>	H	β-acetyloxy	α-acetyloxy	β-CH <sub>3</sub>	β-acetyloxy	H	α-OH	epoxide	α-acetyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>
20) 7-epibaccatin III	α-OH	CH <sub>3</sub>	H	β-acetyloxy	=O	β-CH <sub>3</sub>	α-OH	H	oxetane	α-acetyloxy	α-benzoyloxy	β-OH	H	CH <sub>3</sub>	CH <sub>3</sub>

20. The process of claim 19, wherein said pro-taxanes are isolated from any species of *Taxus*.

21. The process of claim 19, wherein said pro-taxanes are isolated from *Taxus canadensis*.

22. The process of claim 19, wherein the bacteria is *Bacillus cereus* ssp. *taxi*.

23. The process of claim 19, wherein the bacteria is *Bacillus megaterium* ssp. *taxi*.

24. The process of claim 19, wherein the bacteria is *Pantoea* sp. BCM 1.

25. The process of claim 19, wherein the bacteria is *Pantoea* sp. BCM 2.

26. The process of claim 19, wherein the bacteria is *Pantoea* sp. BCM 3.

27. The process of claim 19, wherein the bacteria is *Bacillus cereus* ssp. BCM 4.

28. The process of claim 19, wherein the bacteria is *Bacillus subtilis* ssp. *taxi*.

29. The process of claim 19, wherein the bacteria is *Bacillus megaterium* ssp. BCM 9.

30. The process of claim 19, wherein the bacteria is *Curtobacterium* sp. BCM 5.

31. A process for isolating taxanes and paclitaxel producing bacteria from a yew tree, which comprises the steps of:

- a) disinfecting the surface of a yew tree plant part with a suitable disinfectant;
- b) cutting and disrupting the disinfected plant part;
- c) leaching out bacteria with an appropriate solvent from the part of step c);
- d) culturing said leached bacteria on a solid growth-supporting nutrient medium for a time sufficient to allow formation of visible isolated colonies; and
- e) screening said isolated colonies for production of taxanes and paclitaxel.

32. The process of claim 31, wherein the disinfectant used is ethanol at concentration varying from 35% to 99%.

33. The process of claim 31, wherein the solvent to leach out bacteria is water.

34. The process of claim 31, wherein the solid growth-supporting nutrient medium for isolating colonies is selected from the group consisting of R2A agar (Difco) and Sabouraud (Difco).

35. The process of claim 31, wherein screening of isolated bacteria for taxanes and paclitaxel production comprises the steps of:

- a) culturing said isolated bacteria in 500 ml of a growth-supporting liquid nutrient medium at temperature ranging from 20 to 35°C with constant shaking until sufficient growth is achieved;
- b) obtaining the culture medium supernatant or the cells and extracting it with appropriate organic solvents selected from methylene chloride and ethyl acetate;
- c) analyzing said organic extracts by a method suitable for detection of taxanes and paclitaxel.

36. The process of claim 35, wherein the method to detect taxanes and paclitaxel in extracts uses HPLC retention times on a hydrophobic chromatographic matrix and ultraviolet spectra to identify taxanes and paclitaxel in comparison with standards.

37. The process of claim 35, wherein the method to detect taxanes and paclitaxel in extracts uses cytotoxicity on cancer cell lines.

38. The process of claim 35, wherein the method to detect taxanes in extracts uses mass spectrometry to identify paclitaxel and taxanes based on molecular weights fragments and/or total molecule.

39. The process of claim 35, wherein the method to detect taxanes and paclitaxel in extracts uses antibodies raised against taxanes and paclitaxel.

40. The process of claim 35, wherein the method to



detect taxanes and paclitaxel uses an *in vitro* assay monitoring interaction of taxanes with tubulin.

41. A process for improving biotransformation of pro-taxanes into taxanes and paclitaxel by taxanes and paclitaxel-producing bacteria comprising the steps of:

- a) culturing bacteria in the presence of a mutagenic agent for a time sufficient to allow mutagenesis; and
- b) selecting said mutants by a change of the phenotype which results in an increased biotransformation of pro-taxanes into taxanes and paclitaxel.

42. The process of claim 41, wherein the mutagenic agent is a chemical agent.

43. The process of claim 41, wherein the chemical agent is daunorubicin or nitrosoguanidine.

44. The process of claim 41, wherein the mutagenic agent is a physical agent selected from gamma radiation or ultraviolet.

45. The process of claim 41, wherein the mutagenic agent is a biological agent.

46. The process of claim 45, wherein the biological agent is a transposon.

47. A method of bacterial biotransformation of pro-taxanes into taxanes and paclitaxel thereof which com-

prises the steps of:

- a) incubating the mutated bacteria obtained according to the process of claim 41 in a nutrient medium, and wherein said incubation is effected in the presence of pro-taxanes for a time sufficient to allow production of taxanes and paclitaxel; and
- b) isolating said produced taxanes and paclitaxel thereof from said culturing medium of step a).

48. The process of claim 43, wherein the bacteria used is *Sphingomonas taxi* D200.

49. The process of claim 43, wherein the bacteria used is *Sphingomonas taxi* D201.

50. The process of claim 43, wherein the mutated bacterium is isolated from a *Taxus* species and is of the genus of *Sphingomonas*, *Bacillus*, *Pantoea* or *Curto-bacterium*.

51. A biologically pure culture of a bacteria of genus *Sphingomonas*, *Bacillus*, *Pantoea* or *Curtobacterium*, which is isolated from *Taxus canadensis*, and wherein said bacteria is characterized by the production of taxane and paclitaxel in a culture media.

52. The biologically pure culture of the bacteria of claim 51, which is *Sphingomonas taxi*, *Bacillus cereus* ssp. *taxi*, *Bacillus megaterium* ssp. *taxi*, *Pantoea* sp. BCM 1, *Pantoea* sp. BCM 2, *Pantoea* sp. BCM 3, *Bacillus cereus* ssp. BCM 4, *Bacillus subtilis* ssp. *taxi*, *Bacil-*

*lus megaterium* ssp. BCM 9 and *Curtobacterium* sp. BCM 5.

53. The biologically pure culture of the bacteria of claim 51, wherein the bacteria is of the genus *Sphingomonas*, belonging to the alpha subdivision of Proteobacteria.


54. The biologically pure culture of the bacteria of claim 51, wherein the bacteria is of the genus *Bacillus* belonging to the low G+C Gram positive bacteria.

55. The biologically pure culture of the bacteria of claim 51, wherein the bacteria is of the genus *Pantoea*, belonging to the gamma subdivision of Proteobacteria.

56. The biologically pure culture of the bacteria of claim 51, wherein the bacteria is of the genus is isolated from a *Taxus* species and is of the genus *Curtobacterium*, belonging to the Actinomycetes order of the firmicutes.


57. The biologically pure culture of the bacteria of claim 51, wherein said bacteria is mutated and is *Sphingomonas taxi* D200 or *Sphingomonas taxi* D201.

1/26




SUBSTITUTE SHEET (RULE 26)



2/26

FIG. 1B

3/26



FIG. 2A

4/26

FIG. 2BFIG. 2C



SUBSTITUTE SHEET (RULE 26)

5/26

FIGURE 3AFIGURE 3B





6/26

FIG. 4AFIG. 4B

SUBSTITUTE SHEET (RULE 26)

7/26



Spectral Table


#	Retention Time	Source	Spectrum Name	Baseline Correct	Searchable	Traceable
1	35.820	B.B. F2 def di		On	Yes	Yes
2	35.703	13-mix stds	7-epi-10-deacetyltaxol	On	Yes	Yes

Spectral Table



#	Start Wvln	End Wvln	Resolution	Smooth	Derivative	Spline	Lambda Max	Maximum Absorbance
1	200	350	2.4	None	None	Off	200.6	0.00501
2	200	350	2.4	None	None	Off	200.6	0.42673

5B

8/26

FIG. 6

9/26



SUBSTITUTE SHEET (RULE 26)

60 ATGATTACGC CAAGCTATTT AGGTGACACT ATAGAATACT CAAGCTATGC ATCCAACGCG  
120 TTGGGAGCTC TCCCATATGG TCGACCCTGCA GCGGCCGCA CTAGTGATTA GAGTTTGATC  
180 CTGGCTCAGA ACGAACGCTG GCGGCATGCC TAACACATGC AAGTCGAACG AGATCTTCGG  
240 ATCTAGTGCG GCACGGGTGC GTAACGCGTG GGAATCTGCC CTTTGGTTTCG GAATAACAGT  
300 TGGAACGAC TGCTAATACC GGATGATGAC GTAAGTCCAA AGATTATCG CCAGAGGATG  
360 AGCCCGCGTA GGATTAGCTA GTTGGTGTGG TAAGAGCGCA CCAAGGCGAC GATCCTTAGC  
420 TGGTCTGAGA GGATGATCAG CCACACTGGG ACTGAGACAC GGCCCAGACT CCTACGGGAG  
480 GCAGCAGTGG GGAATATTGG ACAATGGCGG AAAGCCTGAT CCAGCAATGC CCGGTGAGTT  
540 GATGAAAGCC TTAGGTTGTT AAAGCTCTTT TACCCGGGAA TGATAATGAC AGTACCGGA  
600 GAATAAGCTC CGGCTAACTC CGTGCCAGCA GCCCGGGTAA TACGGAAGGA GCTAGCGTTG  
660 TTCGGAATTA CTGGGCGTAA AGCGCACGTA GCGGCTTTG TAAGTTAGAG GTGAAAGCCT  
720 GGAGCTCAAC TCCAGAAATTG CCTTTAAGAC TGCATCGCTT GAATCCAGGA GAGGTGAGTG  
780 GAATTCCGAG TGTAGAGGTG AAATTTCGTAG ATATTTCGGAA GAACACCAGT GCGGAAGGCG  
840 GCTCACTGGA CTGGTATTGA CGCTGAGGTG CGAAAGCGTG GGGAGCAAAC AGGATTAGAT  
900 ACCCTGGTAG TCCACGCCGT AAACGATGAT AACTAGCTGT CCGGGGACTT GGTCTTTGGG  
960 TGGCGCAGCT AACGCATTAA GTTATCCGCC TGGGAGTAC GCGCGCAAGG TTAAAACTCA  
1020 AATGAATTGA CGGGGGCCCTG CACAAGCAGT GAGCATGTG GGTAAATTCTG AAGCAACGCG  
1080 CAGAACCTTA CCAGCGTTTG ACATGTCCCG ACGATTTCTG GAGACAGATC TCTTCCCTTC  
1140 GGGGACTGGA ACGCAGGTGC TGCATGGCTG TCGTCAGCTC GTGTCGTGAG ATGTTGGGTT  
1200 AAGTCCCACA ACGAGCGCAA CCCTCGCCTT TAGTTACCAT CATTAGTTG GGGACTCTAA  
1260 AGGAACCGCC GGTGATAAGC CCGAGGAAGG TGGGATGAC GTCAAGTCCT CATGGCCCTT  
1320 ACGCGCTGGG CTACACACGT GCTACAATGG CCGTGACAGT GGGCAGCAA CTCCGAGAG  
1380 TGCGCTAATC TCCAAAAGCC GTCTCAGTTC GGATTGTTCT CTGCAACTCG AGAGCATGAA  
1440 GGCGGAATCG CTAGTAATCG CCGATCAGCA TGCCGCGGTG AATACGTTCC CAGGCCTTGT  
1500 ACACACCGCC CGTCACACCA TGGGAGTTGG GTTCACCCGA AGGCGTTGCG CTAACTCGTA  
1556 AGAGAGGCAG GCGACCACGG TGGGCTTAGC GACTGGGGTG AAGTCGTAAC AAGGTA

60	AGAGTTTGAT	CATGGCTCAG	GATGAACGCT	GGCGCGGTGC	CTAATACATG	CAAGTCGAGC
120	GAATGGATTA	AGAGCTTGCT	CTTATGAAGT	TAGCGGCCGA	CGGTGAGTA	ACACGTGGGT
180	AACCTACCCA	TAAGACTGGG	ATAACTCCGG	GAAACCGGG	CTAATACCGG	ATAATATTTT
240	GAACTGCATA	GTTTCGAAAT	GAAAGCGGC	TTCGGCTGTC	ACTTATGGAT	GGACCCGCGT
300	CGCATTAGCT	AGTTGGTGAG	GTAAACGGCTC	ACCAAGGCGA	CGATGCGTAG	CCGACCTGAG
360	AGGGTGATCG	GCCACACTGG	GACTGAGACA	CGGCCCAGAC	TCCCTACGGA	GGCAGCAGTA
420	GGGAATCTTC	CGCAATGGAC	GAAAGTCTGA	CGGAGCAACG	CCGCGTGAGT	GATGAAGGCT
480	TTCGGGTCGT	AAAACCTCTGT	TGTTAGGGAA	GAAACAAGTGC	TAGTTGAATA	AGCTGGCACC
540	TTGACGGTAC	CTAACCCAGAA	AGCCACGGCT	AAC TACGTGC	CAGCAGCCCG	CGGTAATACG
600	TAGGTGGCAA	GCGTTATCCG	GAATTATTGG	GCGTAAAGCG	CGCGCAGGTG	GTTTCTTAAG
660	TCTGATGTGA	AAGCCACCGG	CTCAACCGTG	GAGGTCATT	GGAACTGGG	AGACTTGAGT
720	GCAGAAGAGG	AAAGTGGAAT	TCCATGTGTA	GCGGTGAAAT	GCGTAGAGAT	ATGGAGGAAC
780	ACCAGTGGCG	AAGGCGACTT	TCTGGTCTGT	AAC TGACACT	GAGGCGCGAA	AGCGTGGGGA
840	GCAACACAGGA	TTAGATACCC	TGGTAGTCCA	CGCCGTAAAC	GATGAGTGCT	AAGTGTTAGA
900	GGGTTTCCGC	CCTTTAGTGC	TGAAGTTAAC	GCATTAAGCA	CTCCGCCCTGG	GGAGTACGGC
960	CGCAAGGCTG	AAACTCAAAG	GAATTGACGG	GGCCCCGCAC	AAGCGGTGGA	GCATGTGGTT
1020	TAAATTCGAAG	CAACGCCAAG	AACCTTACCA	GGTCTTGACA	TCCTCTGAAA	ACTCTAGAGA
1080	TAGAGCTTCT	CCTTCGGGAG	CAGAGTGACA	GGTGGTG CAT	GGTTGTCgTC	AGCTCGTGTC
1140	GTGAGATGTT	GGGTTAAGTC	CCGCAACGAG	CGCAACCCCTT	GATCTTAGTT	GCCATCATTA
1200	AGTTGGGCAC	TCTAAGGTGA	CTGCCGGTGA	CAAACCGGAG	GAAGGTGGGG	ATGACGTCAA
1260	ATCATCATGC	CCCTTATGAC	CTGGGCTACA	CACGTGCCAC	AATGGACGGT	ACAAAGAGCT
1320	GCAAGACCGC	GAGGTGGAGC	TAACTTCATA	AAACCGTTCT	CAGTTCGGAT	TGTAGGCTGC
1380	AACTCGCCTA	CATGAAGCTG	GAATCGCTAG	TAATCGCGGA	TCAGCATGCC	GCGGTGAATA
1440	CGTCCCGGG	CCTTGTAAC	ACCGCCCGTC	ACACCACGAG	AGTTTGTAAC	ACCCGAAGTC
1500	GGTGGGGTAA	CCTTTATGGA	GCCAGCCGCC	TAAGGTGGGA	CAGATGATTG	GGGTGAAGTC
1512	GTAACAAGGT	AA				



12/26

60 AGAGTTTGAT CCTGGCTCAG GATGAACGCT GGCGGCGTGC CTAATACATG CAAGTCGAGC  
120 GAAC TGATTA GAAGCTTGCT TCTATGACGT TAGCGGCGGA CCGGTGAGTA ACACGTGGGC  
180 AACCTGCCCTG TAAGACTGGG ATAACCTCGG GAAACCGAAG CTAATACCGG ATAGGATCTT  
240 CTCCCTTCATG GGGGATGATT GAAAGATGGT TTCGGCTATC ACTTACAGAT GGGCCCGCGG  
300 TGCAATTAGCT AGTTGGTGAG GTAACGGCTC GACTGAGACA CGGCCCCAAC TCCTACGGGA GGCAGCAGTA  
360 AGGGTGATCG GCCACACTGG CGCAATGGAC GAAAGTCTGA CCGAGCAACG CCGCGTGAGT GATGAAGGCT  
420 GGGAATCTTC CGCAATGGAC AAAACTCTGT TGTnGGGA ACTACTTTCC AGCAGCCGCG GTAATACnTA  
480 TTCGGGTCCT AAAACTCTGT TGTnGGGA ACTACTTTCC ACTACTTTCC GCGGCGCCAGG CGTTGTTCTT  
540 TGACGGTACC TAACCCAGAA CGTTATCCGG AATTATTGGG GGCTCAACCG TGGAGGGTCA TTGGAACCTG GGGAACTTGA  
600 TGTnGGCAAG CGTTATCCGG GAAAGCCCAC GAAAGCGGC GATnAGATAC CACGCCGTAA ACGCATTAAG CACTCCGCCT GGGGAGTACG  
660 AGTCTGATGT GAAAGCGGC GAAAGCGGC GATnAGATAC CACGCCGTAA ACGCATTAAG CACTCCGCCT GGGGAGTACG  
720 GTGCAGAAGA GAAAGCGGC GATnAGATAC CACGCCGTAA ACGCATTAAG CACTCCGCCT GGGGAGTACG  
780 ACACCAGTGG CGAAGCGGC GATnAGATAC CACGCCGTAA ACGCATTAAG CACTCCGCCT GGGGAGTACG  
840 GAGCAAAACAG GATnAGATAC CACGCCGTAA ACGCATTAAG CACTCCGCCT GGGGAGTACG  
900 GAGGGTTTCC GCCCTTTAGT GCTGCAGCTA AGGAATTGAC GGGGGCCCCG CACAGCGGTG GAGCATGTGG  
960 GTCGCAAGAC TGAAACTCAA AGGAATTGAC GGGGGCCCCG CACAGCGGTG GAGCATGTGG  
1020 TTTAATTCTGA AGCAACGCCA AYAACCTTAC CAGGTCTTGA CATCCTCTGA CAACCTCTAGA  
1080 GATAGAGCGT TCCCCTTCGG GGGACAGAGT GACAGGTGGT GCATGGTTGT CGTCAGCTCG  
1140 TGTCTGTGAGA TgTTGGGtTA AGTCCCAGCA AGGCGCAAC CGAGCGCAAC CTTTGATCTT AGTTGCCAGC  
1200 ATTTAGTTGG GCACCTCTAAG GTGACTGCCG GTGACAAACC GTGACAAACC GGAGGAAGGT GGGGATGACG  
1260 TCAAATCATC ATGCCCTTA TGACCTGGGC TACACACGTG CTACAATGGA TGGTACAAAG  
1320 GGCTGCAAGA CCGCGAGGTC AAGCCAATCC CATAAAACCA TTCTCAGTTC GGATTGTAGG  
1380 CTGCAACTCG CCTACATGAA GCTGGAATCG CTAGTAATCG CCGATCAGCA TGCCGCGGTG  
1440 AATACGTTCC CCGGCCCTTGT ACACACCGCC CGTCACACCA CGAGAGTTG TAACACCCGA  
1500 AGTCGGTGGA GTAAACCGTAA GGAGCTAGCC GCCTAAGGTG GGACAGATGA TTGGGGTGAA  
1515 GTCGTAACAA GTAA

SUBSTITUTE SHEET (RULE 26)





TCGGGACGGT CAGCACACGA GGAGCTTGCT CCTTGGGTGA CGAGTGGCGG ACGGGTGAGT 60  
AATGTCGGG GATCTGCCCG ATAnAGGGG ATAACCACTG GAAACGGTGG CTAATAACCGC 120  
ATAACGTCGC AAGACCAAAG AGGGGACCT TCGGGCCTCT CACTATCGGA TGAACCCAGA 180  
TGGGATTAGC TAGTAnGCGG GGTATGGCC CACCTAGGCG ACGATCCCTA nCTGGTCTGA 240  
GAGGATGACC AGCCACACTG GAAC TGAGAC ACGGTCCAnA CTCCTACGGG AGGCAGCAGT 300  
GGGGAATATT GCACAATGGG CGCAAGCCTG ATGCAGCCAT GCCGCGTGTA TGAAGAAGGC 360  
CTTCGGGTG TAAAGTACTT TCAGCGGGA GGAAGCGGAT GCGGTAAATA ACCCTGTCTGA 420  
TTGACGTTCC CCGCAnGAAG AAGCACCGG TAACTCCGTG CCAGCAGCCG CCGTAATACC 480  
GGAGGGTGCA AGCGTTAATC CGGAATTACT GGGCGTAAAG CGCACGCAGG CGGTCTGTTA 540  
AGTCAGATGT GAAATCCCCG GGCTTAACCT GGAACCTGCA TTTGAAACTG GCAGGCTTGA 600  
GTCTTGTAGA GGGGGTAGA ATTCCAGGTG TAGCGGTGAA ATGCCGTAGAG ATCTGGAGGA 660  
ATACCGGTGG CGAAGCGGC GATTAGATAC CCTTGGACA AAGACTGACG CTCAGGTGCG AAAGCGTGGG 720  
GAGCAACAG GATTAGATAC CCTTGGACA AAGACTGACG CTCAGGTGCG AAAGCGTGGG 780  
GTCCCTTGA GGAGTGGCTT CCGGAGCTAA CGCGTTAAGT CACGCCGCTG CGAGGTACGG 840  
CCGCAAGGTT AAAATTCAAA TGAATTGACG GGGGCCCGCA CAAGCGGTGG AGCATGTGGT 900  
TTAATTTCGAT GCAACGCGAA GAACCTTACC TACTCTTGAC ATCCAGCGAA CTTGCCAGAG 960  
ATGGATTGGT GCCTTCGGGA ACGCTGAGAC ACGTGCTGCA TGGCTGTCTG CAGCTCGTGT 1020  
TGTGAAATGT TGGGTTAAGT CCGCAACGA GCGCAACCTT TATCCTTTGT TGCCAGCGAT 1080  
TCGGTCGGGA ACTCAAAGGA GACTGCCGGT GATAAACCGG AGGAAGGTGG GGATGACGTC 1140  
AAGTCATCAT GGCCCTTACG AGTAGGGCTA CACACGTGCT ACAATGGCGC ATACAAAGAG 1200  
AAGCGACCTC GCGAGAGCAA GCGGACCTCA CAAAGTGCGT CGTAGTCCGG ATCGGAGTCT 1260  
GCAACTCGAC TCCGTGAAGT CGGAATCGCT AGTAATCGTG GATCAGAATG CCACGGTGAA 1320  
TACGTTCCCG GGCCTTGTA ACACCGCCCG TCACACCATG GGAGTGGGT GCAAAAAGAG 1380  
TAGGTAGCTT AACCTTCGGG AGGGCGCTTA CCACTTTGTG ATTCATGACT GGGGTGAAGT 1440  
CGTAACAAGT A

FIG. 1

AGAGTTTGAT CCTGGCTCAG GATGAACGCT GCGGCGGTGC CTAATACATG CAAGTCGAGC 60  
GAATGGATTG AGAGCTTGCT CTCAAGAAGT TAGCGGCGGA CGGTGAGTA ACACGTGGGT 120  
AACCTGCCCA TAAGACTGGG ATAACTCCGG GAAACCGGG CTAATACCGG ATAACATTTT 180  
GAACTGCATG GTTCGAAATT GAAAGCGGC GAAAGGCTC ACCAAGGCAA CGATGCGTAG CCGACCTGAG 240  
CGCATTAGCT AGTTGGTGAG GCCACACTGG GACTGAGACA CGGCCCAGAC TCCTACGGGA GGCAGCAGTA 300  
AGGGTGATCG GCGAATGGAC CGCAATGGAC GAAAGTCTGA CGGAGCAACG CCGCGTGAGT GATGAAGGCT 360  
GGGAATCTTC AAAACTCTGT TGTAGGGAA AGCCACGGCT AACTACGTGC GAACAAGTGC TAGTTGAATA AGCTGGCACC 420  
TTCGGGTCGT CTAACCAGAA CGTTATCCGG AATTATTGGG CGTAAAGCGC CAGCAGCCGC GGTAATACGT 480  
TTGACGGTAC AGCCACCGC AAGTGGAAAT CTGGTCTGTA ACTGACACTG CGGTGAAATG CGTAGAGATA TGGAGGAACA 540  
AGGTGGCAAG CGTTATCCGG AGCCACCGC AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 600  
CTGATGTGAA AAGTGGAAAT TAGATACCCT AGGCGACTTT CTTTAGTGCT CTTTAGTGCT GCTCTGAAAA CCCTAGAGAT 660  
CAGAAGAGGA AAGTGGAAAT TAGATACCCT AGGCGACTTT CTTTAGTGCT CTTTAGTGCT GCTCTGAAAA CCCTAGAGAT 720  
CCAGTGGCGA AAGTGGAAAT TAGATACCCT AGGCGACTTT CTTTAGTGCT CTTTAGTGCT GCTCTGAAAA CCCTAGAGAT 780  
CAAACAGGAT TAGATACCCT AGGCGACTTT CTTTAGTGCT CTTTAGTGCT GCTCTGAAAA CCCTAGAGAT 840  
GGTTCCGCC CTTTAGTGCT AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 900  
GCAAGGCTGA AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 960  
AATTCGAAGC AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 1020  
AGGGCTTCTC CTTTAGTGCT AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 1080  
TGAGATGTTG CTTTAGTGCT AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 1140  
GTTGGGCACT CTTTAGTGCT AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 1200  
TCATCATGYC CTTTAGTGCT AACTCAAAGG AATTGACGGG GCGGTGAGT TCCGCCCTGGG GAGTACGGCC 1260  
CAAGACCGCG AGGTGGAGCT AATCTCATAA AACCGTTCTC AGTTCrGATT GTAGGCTGCA 1320  
ACTCGCCTAC ATGAAGCTGG AATCGCTAGT AATCGCGGAT CAGCATGCCG CCGTGAATAC 1380  
GTTCCCGGGC CTTGTACACA CCGCCCGTCA CACCACGAGA GTTTGTAAACA CCCGAAGTCG 1440  
GTGGGGTAAC CTTTTTGGAG CCAGCCGCCT AAGGTGGGAC AGATGATTGG GGTGAAGTCG 1500  
TAACAAGGTA GCC 1513

AGAGTTTGAT CCTGGCTCAG GACGAACGCT GCGGCGGTGC CTAATACATG CAAGTCGAGC 60  
GAACAGATGG GAGCTTGCTC CCTGATGTTA GCGGCGGACG GGTGAGTAAC ACGTGGGTAA 120  
CCTGCCCTGTA AGACTGGGAT AACTCCGGGA AATCGGGGCT AATACCGGAT GGTGTTTGA 180  
ACCGCATGGT TCAGACATAA AAGGTGGCTT CGGCTACCAC TTACAGATGG ACCCGCGCG 240  
CATTAGCTAG TTGGTGAGGT AACGGCTCAC CAAGGCGACG ATGCGTAGCC GACCTGAGAG 300  
GGTGATCGGC CACACTGGGA CTGAGACACG GCCCAGACTC CTACGGGAGG CAGCAGTAGG 360  
GAATCTTCCG CAATGGACGA AAGTCTGACG GAGCAACGCC GCGTGAGTGA TGAAGGTTTT 420  
CGGATCGTAA AGCTCTGTTG TTAGGGAAGA ACAAGTGCCG TTCAAATAGG GCGGCACCTT 480  
GACGGTACCT AACCAGAAAG CCACGGCTAA CTACGTGCCA GCAGCCGCGG TAATACGTAG 540  
GTGGCAAGCG TTGTCCGGAA TTATTGGCGG TAAAGGGCTC GCAGGCGGTT TCTTAAGTCT 600  
GATGTGAAAG CCCCCTGGTC AACGGGGGAG GGTCAATTGA AACTGGGAA CTTGAGTGCA 660  
GAAGAGGAGA GTGGAATTCC ACGTGTAGCG GTGAAATGCG TAGAGATGTG GAGGAACACC 720  
AGTGCGGAAG GCGACTCTCT GGTCTGTAAC TGACGCTGAG GAGCGAAAGC GTGGGGAGCG 780  
AACAGGATTA GATACCCTGG TAGTCCACGC CGTAAACGAT GAGTGCTAAG TGTTAGGGGG 840  
TTTCCGCCCC TTAGTGCTGC AGCTAACGCA TTAGCGGGG CCCGCACAAG CTTGACATCC 900  
AAGACTGAAA CTCAAAGGAA TTGACGGGG CTTGACATCC CTTGACATCC TCTGACAATC 960  
TTCGAAGCAA CGCGAAGAAC CTTACCAGGT CTTGACATCC CTTGACATCC TCTGACAATC 1020  
GACGTCCCCC TCGGGGGCAG AGTGACAGGT AGTGACAGGT GGTGCATGGT TGTCGTCAGC TCGTGTCTGT 1080  
AGATGTTGGG TTAAGTCCC GCAACGAGCGC CAACGAGCGC AACCCCTTGAT CTTAGTTGCC AGCATTCAGT 1140  
TGGGCACTCT AAGGTGACTG CCGGTGACAA ACCGGAGGAA GGTGGGGATG ACGTCAAATC 1200  
ATCATGCCCC TTATGACCTG GGCTACACAC GTGCTACAAT GTGCAGAACA AAGGCAGCG 1260  
AAACCGCGAG GTTAAGCCAA TCCCACAAAT CTGTTCTCAG TTCGGATCCG AGTCTGCAAC 1320  
TCGACTGCGT GAAGCTGGAA TCGCTAGTAA TCGCGGATCA GCATGCCCGG GTGAATACGT 1380  
TCCCCGGCCT TGTACACACC GCCCGTCACA CCACGAGAGT TTGTAACACC CGAAGTCGGT 1440  
GAGGTAACCT TTATGGAGCC AGCCGCCGAA GGTGGGACAG ATGATTGGGG TGAAGTCGTA 1500  
ACAAGGTAAAC CG 1512



GCCTAACAGT GCGAGTCGGA CCGTAGCACA GAGGAGCTTG CTC TTCGGGT GACGAGTGGC 60  
GGACGGGTGA GTAATGTCTG GGGATCTGCC CGATGGAGGG GGATAACCAC TGGAAACGGT 120  
GGCTAATACC GCATAATGTC GCAAGACCAA AGTGGGGGAC CTTCCGGCCT CACACCATCG 180  
GATGAACCCA GATGGGATTA GCTAGTAGGT GGGTAACGG CTCACCTAGG CGACGATCCC 240  
TAGCTGGTCT GAGAGGATGA CCAGCCACAC TGGAACTGAG ACACGGTCCA GACTCCTACG 300  
GGAGGCAGCA GTGGGGAATA TTGCACAATG GCGCAAGCC TGATGCAGCC ATGCCGCGTG 360  
TATGAAGAAG GCCTTCGGGT TGTAAGAATAC TTTCACCGG GAGGAAGGG ACGAGGTTAA 420  
TAACCCCGTT CATTGACGTT ACCCGCAGAA GAAGCACCGG CTAACTCCGT GCCAGCAGCC 480  
GCGGTAATAC GGAGGGTGCA AGCGTTAATC GGAATTACTG GCGTAAAGC GCACGCAGGC 540  
GGTCTGTTAA GTCAGATGTG AAATCCCCGG GCTTAACCTG GGAACCTGCAT TTGAAACTGG 600  
CAGGCTTGAG TCTTGTAGAG GGGGTAGAA TTCCAGGTGT AGCGGTGAAA TGCGTAGAGA 660  
TCTGGAGGAA TACCGGTGGC GAAGCGGCC CCCTGGACAA AGACTGACGT TCAGGTGCGA 720  
AAGCGTGGG AGCAAACAGG ATTAGATACC CTGGTAGTCC ACGCCGTAAA CGATGTCGAC 780  
TTGGAGGCTG TGAGCATGAT TCGTGGCTTC CCGAGCTAAC GCGTTAAGTC GACCGCCTGG 840  
GGAGTACGGC CGCAAGGTTA AAATTCAAAT GAATTGACGG GGGCCCGCAC AAGCGGTGGA 900  
GCATGTGGTT TAATTcGATG CAACGCCAAG AACCTTACCT GYTCTTGACA TCCACGGAAT 960  
TYGGCAGAGA TGCCTTAGTG CCTTCGGGAA CCGTGAGACA CCGTGCTGCAT GGCTGTCGTC 1020  
AGCTCGTGTT GTGAAATGTT GGGTTAAGTC CCGCAACGAG CCGCAACCCTT ATCCTTTGTT 1080  
GCCAGCGATT CCGTCGGGAA CTCAAAGGAG ACTGCCGGTG ATAAACC GGAAGGTGGG 1140  
GATGACGTCA AGTCATCATG GCCCTTACGA GCAGGGCTAC ACACGTGCTA CAATGGCGCA 1200  
TACAAAGAGA AGCGACCCTG CGAGAGCAAG CCGACCTCAC AAAGTGCGTC GTAGTCCGGA 1260  
TCGGAGTCTG CAACTCGACT CCGTGAAGTC GGAATCGCTA GTAATCGTGG ATCAGAAATGC 1320  
CACGGTGAAT ACGTTCCCGG GCCTTGTAACA CACCGCCCGT CACACCATGG GAGTGGGTG 1380  
CAAAAGAAGT AGGTAGCTTA ACCTTCCGGA GGGCGCTTAC CACTTTGTGA TTCATGACTG 1440  
GGGTGAAGTC GTAACAAG 1458

60	AGTGGAGTC	GGACGGTAGC	ACAGAGAGCT	TGCTCTTGGG	TGACGAGTGG	CGGACGGGTG
120	AGTAATGTCT	GGGATCTGC	CCGATAGAGG	GGGATAACCA	CTGGAAACGG	TGGCTAATAC
180	CGCATAACGT	CGCAAGACCA	AAGAGGGGA	CCTTCGGGC	TCTCACTATC	GGATGAACCC
240	AGATGGGATT	AGCTAGTAGG	CGGGTAATG	GCCACCTAG	GCGACGATCC	CTAGCTGGTC
300	TGAGAGGATG	ACCAGCCACA	CTGGAAC TGA	GACACGGTCC	AGACTCCTAC	GGGAGGCAGC
360	AGTGGGGAAT	ATTGCACAAT	GGGCGCAAGC	CTGATGCAGC	CATGCCCGCT	GTATGAAGAA
420	GGCCTTCGGG	TTGTAAAGTA	CTTTCAGCGG	GGAGGAAGGC	GATGCCGTTA	ATAACCTGT
480	CGATTGACGT	TACCCGCAGA	AGAAGCACCG	GCTAACTCCG	TGCCAGCAGC	CGCGGTAATA
540	CGGAGGGTGC	AAGCGTTAAT	CGGAATTACT	GGGCGTAAAG	CGCACGCAGG	CGGTCTGTTA
600	AGTCAGATGT	GAAATCCCCG	GGCTTAACCT	GGGAAC TGA	TTTGAAACTG	GCAGGCTTGA
660	GTCCTGTAGA	CGGGGGTAGA	ATTCAGGTG	TAGCGGTGAA	ATGCCGTAGAG	ATCTGGAGGA
720	ATACCGGTGG	CGAAGGCGGC	CCCCTGACA	AAGACTGACG	CTCAGGTGCG	AAAGCGTGGG
780	GAGCAACACAG	GATTAGATAC	CCTGGTAGTC	CACGCCGTAA	ACGATGTCGA	CTTGGAGGTT
840	GTTCCCTTGA	GGAGTGGCTT	CCGGAGCTAA	CGCGTTAAGT	CGACCGCCTG	GGGAGTACGG
900	CCGCAAGGTT	AAAATTCAAA	TGAATTGACG	GGGGCCCGCA	CAAGCGGTGG	AGCATGTGGT
960	TTAATTTCGAT	GCAACGCGAA	GAACCTTACC	TACTCTTGAC	ATCCACGGAA	TTYGGCAGAG
1020	ATGCTTTGGT	GCCTTCGGGA	ACCCTGAGAC	AGGTGCTGCA	TGGCTGTCTG	CAGCTCGTGT
1080	TGTGAAATGT	TGGGTTAAGT	CCCGCAACGA	GCGCAACCCT	TATCCTTTGT	TGCCAGCGAT
1140	TCGGTCGGGA	ACTCAAAGGA	GACTGCCGGT	GATAAACCGG	AGGAAGGTGG	GGATGACGTC
1200	AAGTCATCAT	GGCCCTTACG	AGTAGGGCTA	CACACGTGCT	ACAATGGCGC	ATACAAAGAG
1260	AAGCGACCTC	GCGAGAGCAA	GCGGACCTCA	CAAAGTGCGT	CGTAGTCCGG	ATCGGAGTCT
1320	GCAACTCGAC	TCCGTGAAGT	CGGAATCGCT	AGTAATCGTG	GATCAGAATG	CCACGGTGAA
1380	TACGTTCCCG	GGCCTTG TAC	ACACCGCCCG	TCACACCATG	GGAGTGGGTT	GCAAAAGAAG
1440	TAGGTAGCTT	AACCTTCGGG	AGGGCGCTTA	CCACTTTGTG	ATTCA TGACT	GGGTGAAGT
1451	CGTAACAAGT	A				

FIG. 1 - BH

60	AGAGTTTGAT	CATGGCTCAG	GATGAACGCT	GGCGGCGTGC	CTAATACATG	CAAGTCGAGC
120	GAACTGATTA	GAAGCTTGCT	TCTATGACGT	TAGCGGCGGA	CGGTGAGTA	ACACGTGGGC
180	AACCTGCCCTG	TAAGACTGGG	ATAACTTCGG	GAAACCGAAG	CTAATACCGG	ATAGGATCTT
240	CTCCTTCATG	GGAGATGATT	GAAAGATGGT	TTCGGCTATC	ACTTACAGAT	GGCCCCCGCG
300	TGCATTAGCT	AGTTGGTGAG	GTAAACGGCC	ACCAAGGCAA	CGATGCATAG	CCGACCCTGAG
360	AGGGTGATCG	GCCACACTGG	GACTGAGACA	CGGCCAGAC	TCCTACGGGA	GGCAGCAGTA
420	GGGAATCTTC	CGCAATGGAC	GAAAGTCTGA	CGGAGCAACG	CCGCGTGAGT	GATGAAGGCT
480	TTCGGGTCTG	AAAACCTCTG	TGTTAGGGAA	GAACAAGTAC	GAGAGTAACT	GCTCGTACCT
540	TGACGGTACC	TAACCCAGAA	GCCACGGCTA	ACTACGTGCC	AGCAGCCCGG	GTAATACGTA
600	GGTGGCAAGC	GTTATCCGGA	ATTATTGGGC	GTAAAGCGCG	CCGCAGGCGG	TTTCTTAAGT
660	CTGATGTGAA	AGCCCACGGC	TCAACCGTgG	GArGGTCATT	GGAAACTGGG	GAACTTGAgt
720	GCAGAAGAGA	AAAGCGGAAT	TCCACGTGTA	GCGGTGAAAT	GCGTAGAGAT	GTGGAGGAAC
780	ACCAGTGGCG	AAGGCGGCTT	TTTGGTCTGT	AACTGACGCT	GAGGCGCGAA	AGCGTGGGGA
840	GCAAAACAGGA	TTAGATACCC	TGGTAGTCCA	CGCCGTAAAC	GATGAGTGCT	AAGTGTTAGA
900	GGGTTTCCGC	CCTTTAGTGC	TGCAGCTAAC	GCATTAAAGCA	CTCCGCCCTGG	GGAGTACGGT
960	CGCAAGACTG	AAACTCAAAG	GAATTGACGG	GGGCCCGCAC	AAGCGGTGGA	GCATGTGGTT
1020	TAATTTCGAAG	CAACGCGAAG	AACCTTACCA	GGTCTTGACA	TCcTCTGACA	ACTCTAGAGA
1080	TAGAGCtTTC	CCcTTCCGGG	GACACAGTGA	CAGGTGGTGC	ATGGTTGTCC	TCAGCTCGTG
1140	TCGTGAGATG	TTGGGTTAAG	TCCCGCAACG	AGCGCAACCC	TTGATCTTAG	TTGCCAGCAT
1200	TTAGTTGGGC	ACTCTAAGGT	GACTGCCGGT	GACAAACCCG	AGGAAGGTGG	GGATGACGTC
1260	AAATCATCAT	GCCCCTTATG	ACCTGGGCTA	CACACGTGCT	ACAATGGATG	GTACAAAGGG
1320	CTGCAAGACC	GCGAGGTCAA	GCCAATCCCA	TAAAACCATT	CTCAGTTCGG	ATTGTAGGCT
1380	GCAACTCGCC	TACATGAAGC	TGGAATCGCT	AGTAATCGCG	GATCAGCATG	CCGCGGTGAA
1440	TACGTTCCCCG	GGCCTTGTA	ACACCGCCCC	TCACACCCAG	AGAGTTTGTA	ACACCCGAAG
1500	TCGGTGGAGT	AACCGTAAGG	AGCTAGCCCC	CTAAGGTGGG	ACAGATGATT	GGGGTGAAGT
1513	CGTAACAAGG	TAG				



CGCTGGGCGG CCTGCTTAAC ACnTGCAAGT CGAACGATGA TGCCCAGCTT GCTGGGTGGA 60  
TTAGTGCGGA ACGGGTGAGT AACACGTGAG TAACCTGCCC CTGACTCTGG GATAAGCGTT 120  
GGAAACGACG TCTAATACTG GATATGATCA CTGGCCGCAT CTGCTGGTGG TGGAAAGATT 180  
TTTTGGTTGG GGATGGACTC GCGGCCATATC AGCTTGTTGG TGAGGTAATG GCTCACCAAG 240  
GCGACGACGG GTAGCCGGCC TGAGAGGGTG ACCGGCCACA CTGGGACTGA GACACGGCCC 300  
AGACTCCTAC GGGAGGCAAC AGTGGGGAAT ATTGCACAAT GGGCGAAAGC CTGATGCAGC 360  
AACGCCGCGT GAGGGATGAC GGCCTTCGGG TTGTAAACCT CTTTTAGTAG GGAAGAAGCG 420  
AAAGTGACGG TACCTGCAGA AAAAGCACCG GCTAACTACG TGCCAGCAGC CGCGGTAATA 480  
CGTAGGGTGC AAGCGTTGTC CGGAATTATT CGGCGTAAAG AGCTCGTAGG CGGTTGTGCG 540  
CGTCTGCTGT GAAATCCCGA GGCTCAACCT CCGGCTTGCA GTGGGTACGG GCAGACTAGA 600  
GTGCGGTAGG GGAGATTGGA ATTCTTGGTG TAGCGGTGGA ATGCGCAGAT ACCAGGAGGA 660  
ACACCGATGG CGAAGGCAGA TCTCTGGGCC GTAAC TGACG CTGAGGAGCG AAAGCATGGG 720  
GAGCGAACAG GATTAGATAC CCTGGTAGTC CATGCCGTAA ACGTTGGCG CTAGATGTAG 780  
GGACCTTTCC ACGGTTTCTG TGTCGTAGCT AACGCATTAA GCGCCCCGCC TGGGGAGTAC 840  
GGCCGCAAGG CTAAAACTCA AAGGAATTGA CCGGGGCCCG CACAAGCGGC GGAGCATGCG 900  
GATTAAATTCG ATGCAACGCG AAGAACCCTA CCAAGGmTTG ACATACACCG GAAACGGCCA 960  
GAGATGGTCG CCCCCTTGTG GTCGGTGTAC AGGTGGTGCA TGGTTGTCGT CAGCTCGTGT 1020  
CGTGAGATGT TGGGTTAAGT CCGGCAACGA AGACTGCCCG GGTCAACTCG GAGGAAGGTG 1080  
TTATGCCGGG GACTCATAGG AGACTGCCCG GTCAACTCG GGTCAATGGCC GGTACAAAGG 1140  
CAAATCATCA TGCCCCCTTAT GTCTTGGGCT TCACGCATGC TACAATGGCC GGTACAAAGG 1200  
GCTGCCGATAC CGTAAGGTGG AGCGAATCCC AAAAAAGCCGG TCTCAGTTCTG GATTGAGGTC 1260  
TGCAACTCGA CCTCATGAAG TCGGAGTCGC TAGTAATCGC AGATCAGCAA CGCTGCCGTG 1320  
AATACGTTCC CGGGCCTTGT ACACACCGCC CGTCAAGTCA TGAAGTCGG TAACACCCGA 1380  
AGCCGGTGGC CTAACCCCTG TGGAAGGAGC CGTCGAAGGT GGGATCGGTG ATTAGGACTA 1440  
AGTCGTAACA AG





20/26


AGAGTTTGAT CATGGCTCAG AACGAACGCT GGCGGCATGC CTAAACATG CAAGTCGAAC 60  
GAGATCTTCG GATCTAGTGG CGCACGGGTG CGTAACGCgT gGGAATCTGC cCTTtGGTTC 120  
GGAATAaCAG TTGGAAACGA CTGCTAATgA CCGGATGATG ACGTAAGTCC AAAGATTAT 180  
CGCCAGAGGA TkAGCCCGCG TAGGATTAGC tAgTTGGTGT gGTaaGAGCG CACCAAGGCG 240  
ACgAtCCTTA GCTGGTCTGA GArGATGAtC GArGATGAtC AGCCACACTG GGACTGAGAC ACGGCCAgA 300  
CTCCTACGGG AGGCAGCAGT AGGCAGAtwTT GGGGAAtwTT GGACAATGGG CGAAAGCCTG ATCCAGCAAT 360  
GCCGCGTGAG TGATGAAGGC CTTAGGGTTG TAAAGCTCTT TTACCCGGGA TGATAATGAC 420  
AGTACC GGGA GAATAAGCTC CGGCTAACTC CGTGCCAGCA CGCGCGGTAA TACGGAGGGA 480  
GCTAGCGTTG TTCGGAATTA CTGGGCGTAA AGCGCACGTA AGCGGCTTTG TAAGTTAGAG 540  
GTGAAAGCCT GGAGCTCAAC GGAGCTCAAC TCCAGAATTG CCTTTAAGAC TGCATCGCTT GAATCCAGGA 600  
GAGGTGAGTG GAATTCCGAG GAATTCCGAG TGTAGAGGTG AAATTCTAG AAATTCTGAA GAACACCAGT 660  
GGCGAAGCG GCTCACTGGA ACCCTGGTAG TCCACGCCGT AACCGATGAT CGCTGAGGTG CGAAAGCGTG GGGAGCAAAAC 720  
AGGATTAGAT ACCCTGGTAG TGGCGCAGCT AACGCATTAA GTTATCCGCC AAACGATGAT AACTAGCTGT CCGGGGACTT 780  
GGTCTTTGGG TGGCGCAGCT AATGAATTGA CGGGGGCCTG CACAAAGCGGT GGAGCATGTG GTTTAATTCTG 900  
AAGCAACCG CAGAACCTTA CAGCGGTTTG CCAGCGTTTG ACATGTCCGG ACATTTCTG GAGACAGATC 960  
TCTTCCCTTC GGGGACTGGA AAGTCCCGCA ACGAGCGCAA CCTCGCCTT TCGTCAGCTC GTGTCGTGAG 1020  
ATGTTGGGTT AAGTCCCGCA AAGTCCCGCA ACGAGCGCAA CCTCGCCTT TAGTTACCAT CATTAGTTG 1080  
GGGACTCTAA AGGAACCGCC AGGAACCGCC GGTGATAAGC GGTGATAAGC CGGTGACAGT GGCAGCAAA 1200  
CATGGCCCTT ACGCGCTGGG ACGCGCTGGG CTACACACGT CTACACACGT GCTCAATGG GTCAAGTCTT 1260  
CTCGCGAGAG TCGGCTAATC TCGGCTAATC TCCAAAAGCC GTCTCAGTTC GGATTGTTCT CTGCAACTCG 1320  
AGAGCATGAA GCGGGAATCG GCGGGAATCG CTAGTAATCG CGGATCAGCA TGCCGCGGTG AATACGTTCC 1380  
CAGGCCCTTG ACACACCGCC ACACACCGCC CGTCACACCA TGGGAGTTGG GTTCACCCGA AGCGTTGCG 1440  
CTAACTCAGC AATGAGAGGC AGGCGACCAC AGGCGACCAC GGTGGGCTTA GCGACTGGGG TGAAGTCGTA 1450  
ACAAGGTAAC

SUBSTITUTE SHEET (RULE 26)

FIG. 1 - BK





22/26

FEI-10A



m/z	Assignment
447	(T-BzOH) <sup>+</sup>
386	(T-H <sub>2</sub> O-BzOH-AcOH) <sup>+</sup>
326	(T-BzOH-2AcOH-H <sub>2</sub> O) <sup>+</sup>
308	(T-BzOH-2AcOH-2H <sub>2</sub> O) <sup>+</sup>

FEI-10B



23/26

FIG. 11AFIG. 11B

24/26




25/26



SUBSTITUTE SHEET (RULE 26)

26/26

FIG. 14C

## SEQUENCE LISTING

<110> BCM Développement Inc.  
 PAGÉ, Michel  
 LANDRY, Nathalie  
 BOISSINOT, Maurice  
 HÉLIE, Marie-Claude  
 HARVEY, Mario  
 GAGNÉ, Martin

<120> BACTERIAL MASS PRODUCTION OF TAXANES AND  
 PACLITAXEL

<130> 12637-2PCT FC/ld

<150> US 08/995,960

<151> 1997-12-22

<160> 13

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1556

<212> DNA

<213> Sphingomonas taxi

<400> 1

atgattacgc	caagctat	ttt	agg	tgacact	atagaata	ctatgc	atccaacgcg	60
ttgggagctc	tcccatat	gg	tcgac	ctgca	ggcggccgca	ctagt	gatta	120
ctggctcaga	acgaacgc	tg	gcggc	atgcc	taacacatgc	aagtcgaacg	agatcttcgg	180
atctagtggc	gcacgggtgc	gtaacgc	gtg	ggaatctgcc	ctttgggttcg	gaataacagt		240
tggaaacgac	tgctaata	acc	ggat	gatgac	gtaagtccaa	agatttatcg	ccagaggatg	300
agccccgcgta	ggattagcta	gttgggtgtg	taagagc	gcga	ccaaggcgac	gac	cttagc	360
tggtctgaga	ggatgatcag	ccacactggg	actgagacac	ggcccagact	cctacgggag			420
gcagcagtgg	ggaatattgg	acaatggg	cg	aaagcctgat	ccagcaatgc	cgcgtgagtt		480
gatgaaagcc	ttaggttg	tt	aaagctcttt	tacccgggaa	tgataatgac	agtaccggga		540
gaataagctc	cggctaactc	cgtgcccagca	gccgcggtaa	tacggaagga	gctagcgttg			600
ttcgggaatta	ctgggcgtaa	agcgcacgta	ggcggc	tttg	taagttagag	gtgaaagcct		660
ggagctcaac	tccagaattg	cctttaagac	tgcatcgctt	gaatccagga	gaggtgagtg			720
gaattccgag	tgtagaggtg	aaattcgtag	atattcggaa	gaacaccagt	ggcgaaggcg			780
gctcactgga	ctggtattga	cgctgaggtg	cgaaagcgtg	gggagcaa	ac	aggattagat		840
accctggtag	tccacgccgt	aaacgatgat	aactagctgt	ccggggactt	ggtctttggg			900
tggcgcagct	aacgcattaa	gttatccgcc	tggggagtag	ggccgcaagg	ttaaaactca			960
aatgaattga	cgggggcctg	cacaagcgg	ggagcatgtg	gtttaattcg	aagcaacgcg			1020
cagaacctta	ccagcgtttg	acatgtccgg	acgatttctg	gagacagatc	tcttcccttc			1080
ggggactgga	acgcaggtgc	tgcatggctg	tcgtcagctc	gtgtcgtgag	atgttgggtt			1140
aagtcccgca	acgagcgcaa	ccctcgcctt	tagttaccat	catttagttg	gggactctaa			1200
aggaaccgcc	ggtgataagc	cggaggaagg	tggggatgac	gtcaagtcct	catggccctt			1260
acgcgctggg	ctacacacgt	gctacaatgg	cggtgacagt	gggcagcaaa	ctcgcgagag			1320
tgcgctaate	tccaaaagcc	gtctcagttc	ggattgttct	ctgcaactcg	agagcatgaa			1380
ggcggaatcg	ctagtaatcg	cggatcagca	tgccgcgggtg	aatacgttcc	caggccttgt			1440
acacaccgcc	cgtcacacca	tgggagttgg	gttcacccga	aggcgttgcg	ctaactcgta			1500
agagaggcag	gcgaccacgg	tgggcttagc	gactgggggtg	aagtcgtaac	aaggta			1556



<210> 2  
 <211> 1512  
 <212> DNA  
 <213> *Bacillus cereus* ssp. *taxi*

<400> 2

agagtttgat	catggctcag	gatgaacgct	ggcggcgtgc	ctaatacatg	caagtcgagc	60
gaatggatta	agagcttgct	cttatgaagt	tagcggcgga	cgggtgagta	acacgtgggt	120
aacctacca	taagactggg	ataactccgg	gaaaccgggg	ctaataccgg	ataatatatt	180
gaactgcata	gttcgaaatt	gaaaggcggc	ttcggctgtc	acttatggat	ggacccgcgt	240
cgcattagct	agttggtgag	gtaacggctc	accaaggcga	cgatgcgtag	ccgacctgag	300
aggggtgatcg	gccacactgg	gactgagaca	cggcccagac	tcctacggga	ggcagcagta	360
gggaatcttc	cgcaatggac	gaaagtctga	cggagcaacg	ccgcgtgagt	gatgaaggct	420
ttcgggtcgt	aaaactctgt	tgtaggggaa	gaacaagtgc	tagttgaata	agctggcacc	480
ttgacggtac	ctaaccagaa	agccacggct	aactacgtgc	cagcagcccg	cggtaatacg	540
taggtggcaa	gcgttatccg	gaattattgg	gcgtaaagcg	cgcgaggtg	gtttcttaag	600
tctgatgtga	aagcccacgg	ctcaaccgtg	gagggtcatt	ggaaactggg	agacttgagt	660
gcagaagagg	aaagtggaat	tccatgtgta	gcggtgaaat	gcgtagagat	atggagggaac	720
accagtggcg	aaggcgactt	tctggtctgt	aactgacact	gaggcgcgaa	agcgtgggga	780
gcaaacagga	ttagataccc	tggtagtcca	cgccgtaaac	gatgagtgct	aagtgttaga	840
gggtttccgc	ccttttagtg	tgaagttaac	gcattaagca	ctccgcctgg	ggagtacggc	900
cgcaaggctg	aaactcaaag	gaattgacgg	gggcccgcac	aagcgggtga	gcatgtgggt	960
taattcgaag	caacgcgaag	aaccttacca	ggtcttgaca	tcctctgaaa	actctagaga	1020
tagagcttct	ccttcgggag	cagagtgaca	ggtggtgcat	ggttgctcgt	agctcgtgtc	1080
gtgagatgtt	gggttaagtc	ccgcaacgag	cgcaaccctt	gatcttagtt	gccatcatta	1140
agttgggcac	tctaagggtga	ctgccgggtga	caaaccggag	gaagggtggg	atgacgtcaa	1200
atcatcatgc	cccttatgac	ctgggctaca	cacgtgccac	aatggacggg	acaaagagct	1260
gcaagaccgc	gaggtggagc	taatctcata	aaaccgttct	cagttcggat	tgtaggctgc	1320
aactcgctta	catgaagctg	gaatcgctag	taatcgcgga	tcagcatgcc	gcggtgaata	1380
cgttcccggg	ccttgtacac	accgcccgtc	acaccacgag	agtttgtaac	accogaagtc	1440
ggtggggtaa	cctttatgga	gccagccgcc	taagggtggga	cagatgattg	gggtgaagtc	1500
gtaacaaggt	aa					1512

<210> 3  
 <211> 1515  
 <212> DNA  
 <213> *Bacillus megaterium* ssp. *taxi*

<400> 3

agagtttgat	cctggctcag	gatgaacgct	ggcggcgtgc	ctaatacatg	caagtcgagc	60
gaactgatta	gaagcttgct	tctatgacgt	tagcggcgga	cgggtgagta	acacgtgggc	120
aacctgcctg	taagactggg	ataacttcgg	gaaaccgaag	ctaataccgg	ataggatctt	180
ctccttcctg	ggggatgatt	gaaagatggt	ttcggctatc	acttacagat	gggcccgcgg	240
tgcattagct	agttggtgag	gtaacggctc	accaaggcaa	cgatgcatag	ccgacctgag	300
aggggtgatcg	gccacactgg	gactgagaca	cggcccacaa	tcctacggga	ggcagcagta	360
gggaatcttc	cgcaatggac	gaaagtctga	cggagcaacg	ccgcgtgagt	gatgaaggct	420
ttcgggtcct	aaaactctgt	tgtaggggaa	gaacaaatac	cagagtaact	gcttggttct	480
tgacgggtacc	taaccagaaa	gccacggcta	actactttcc	agcagccgcg	gtaatacnta	540
tgtaggcaag	cggttatccg	aattattggg	gcgttaaagc	gcgcgccagg	cggtttctta	600
agtctgatgt	gaaagcccac	ggctcaaccg	tggagggtca	ttggaaactg	gggaacttga	660
gtgcagaaga	gaaaagcggg	attccacgtg	tagcggtgaa	atgcgtagag	atgtggagga	720
acaccagtgg	cgaaggcggc	tttttggtct	gtaactgacg	ctgaggcgcg	aaagcgtggg	780
gagcaaacag	gattagatac	cctggtagtc	cacgccgtaa	acgatgagtg	ctaagtgtta	840
gagggtttcc	gccctttagt	gctgcagcta	acgcattaag	cactccgcct	ggggagtacg	900
gtcgcaagac	tgaaactcaa	aggaattgac	gggggcccgc	acaagcgggt	gagcatgtgg	960
tttaattcga	agcaacgcga	araaccttac	caggctctga	catcctctga	caactctaga	1020

gatagagcgt	tccccttcgg	gggacagagt	gacaggtggt	gcatgggtgt	cgtcagctcg	1080
tgctcgtgaga	tggtgggtta	agtcccgcaa	cgagcgcaac	ccttgatctt	agttgccagc	1140
atttagttgg	gcactctaag	gtgactgccg	gtgacaaacc	ggaggaaggt	ggggatgacg	1200
tcaaatcatc	atgcccctta	tgacctgggc	tacacacgtg	ctacaatgga	tggtacaaag	1260
ggctgcaaga	ccgcgaggtc	aagccaatcc	cataaaacca	ttctcagttc	ggattgtagg	1320
ctgcaactcg	cctacatgaa	gctggaatcg	ctagtaatcg	cggatcagca	tgccgcggtg	1380
aatacgttcc	cgggccttgt	acacaccgcc	cgtcacacca	cgagagtttg	taacacccga	1440
agtcggtgga	gtaaccgtaa	ggagctagcc	gcctaagggtg	ggacagatga	ttgggggtgaa	1500
gtcgtaacaa	ggtaa					1515

&lt;210&gt; 4

&lt;211&gt; 1451

&lt;212&gt; DNA

&lt;213&gt; Pantoea sp. BCM 1

&lt;400&gt; 4

tcgggacggt	cagcacacga	ggagcttgct	ccttgggtga	cgagtggcgg	acgggtgagt	60
aatgtctggg	gatctgcccg	atanaggggg	ataaccactg	gaaacggtgg	ctaataccgc	120
ataacgtcgc	aagaccaaag	agggggacct	tcgggcctct	cactatcgga	tgaaccaga	180
tgggattagc	tagtangcgg	ggtaatggcc	cacctaggcg	acgatcccta	nctggtctga	240
gaggatgacc	agccacactg	gaactgagac	acggtccana	ctcctacggg	aggcagcagt	300
ggggaatatt	gcacaatggg	cgcaagcctg	atgcagccat	gccgcgtgta	tgaagaaggc	360
cttcgggttg	taaagtactt	tcagcgggga	ggaaggcgat	gcggttaata	accctgtcga	420
ttgacgttcc	ccgcangaag	aagcaccggc	taactccgtg	ccagcagccg	cggtaatacc	480
ggaggggtgca	agcgttaatc	cggaattact	gggcgtaaa	cgcacgcagg	cggctctgta	540
agtcagatgt	gaaatccccg	ggcttaacct	gggaactgca	tttgaaactg	gcaggcttga	600
gtcttgtaga	ggggggtaga	attccagggtg	tagcggtgaa	atgcgtagag	atctggagga	660
ataccggtgg	cgaaggcggc	cccttgga	aagactgacg	ctcaggtgcg	aaagcgtggg	720
gagcaaacag	gattagatac	cctggtagtc	cacgccgtaa	acgatgtcga	cttgagggtt	780
gttcccttga	ggagtggctt	ccggagctaa	cgcgttaagt	cgaccgcctg	gggagtacgg	840
ccgcaagggt	aaaattcaaa	tgaattgacg	ggggcccgc	caagcgggtg	agcatgtggt	900
ttaattcgat	gcaacgcgaa	gaaccttacc	tactcttgac	atccagcgaa	cttgccagag	960
atggattggt	gccttcggga	acgctgagac	aggtgctgca	tggctgtcgt	cagctcgtgt	1020
tgtgaaatgt	tgggttaagt	cccgcaacga	gcgcaaccct	tatcctttgt	tgccagcgat	1080
tcggtcggga	actcaaagga	gactgccggt	gataaaccgg	aggaagggtg	ggatgacgtc	1140
aagtcatcat	ggcccttacg	agtagggcta	cacacgtgct	acaatggcgc	atacaaagag	1200
aagcgacctc	gcgagagcaa	gcggacctca	caaagtgcgt	cgtagtccgg	atcggagtct	1260
gcaactcgac	tccgtgaagt	cggaatcgct	agtaatcggt	gatcagaatg	ccacgggtgaa	1320
tacgttcccc	ggccttgtac	acaccgccc	tcacaccatg	ggagtgggtt	gcaaaagaag	1380
taggtagctt	aaccttcggg	agggcgctta	ccactttgtg	attcatgact	gggggtgaagt	1440
cgtaacaagt	a					1451

&lt;210&gt; 5

&lt;211&gt; 1513

&lt;212&gt; DNA

&lt;213&gt; Bacillus cereus ssp. BCM 4

&lt;400&gt; 5

agagtttgat	cctggctcag	gatgaacgct	ggcggcgtgc	ctaatacatg	caagtcgagc	60
gaatggattg	agagcttgct	ctcaagaagt	tagcggcgga	cgggtgagta	acacgtgggt	120
aacctgcca	taagactggg	ataactccgg	gaaaccgggg	ctaataccgg	ataacatttt	180
gaactgcatg	gttcgaaatt	gaaaggcggc	ttcggctgtc	acttatggat	ggacccgcgt	240
cgcattagct	agttgggtgag	gtaacggctc	accaaggcaa	cgatgcgtag	ccgacctgag	300
aggggtgatcg	gccacactgg	gactgagaca	cggcccagac	tcctacggga	ggcagcagta	360
gggaatcttc	cgcaatggac	gaaagtctga	cggagcaacg	ccgcgtgagt	gatgaaggct	420

ttcggggtcgt	aaaactctgt	tgtaggggaa	gaacaagtgc	tagttgaata	agctggcacc	480
ttgacggtac	ctaaccagaa	agccacggct	aactacgtgc	cagcagccgc	ggtaatacgt	540
aggtggcaag	cggtatccgg	aattattggg	cgtaaagcgc	gcgcaggtgg	tttcttaggt	600
ctgatgtgaa	agcccacggc	tcaaccgtgg	aggggtcattg	gaaactggga	gacttgagtg	660
cagaagagga	aagtggaatt	ccatgtgtag	cggtgaaatg	cgtagagata	tggaggaaca	720
ccagtggcga	aggcgacttt	ctggtctgta	actgacactg	aggcgcgaaa	gcgtggggag	780
caaacaggat	tagataccct	ggtagtccac	gccgtaaacg	atgagtgcta	agtgttagag	840
ggtttccgcc	ctttagtgtc	gaagttaacg	cattaagcac	tccgcctggg	gagtacggcc	900
gcaaggctga	aactcaaagg	aattgacggg	ggcccgcaca	agcgggtggag	catgtggttt	960
aattcgaagc	aacgcgaaga	accttaccag	gtcttgacat	cctctgaaaa	ccctagagat	1020
agggcttctc	cttcggggagc	agagtgcacg	gtgggtgcatg	gttgctcgtca	gctcgtgtcg	1080
tgagatgttg	ggttaagtcc	cgcaacgagc	gcaacccttg	atcttagttg	ccatcattaa	1140
gttgggcact	ctaagggtgac	tgccgggtgac	aaaccggagg	aaggtgggga	tgacgtcaaa	1200
tcatcatgyc	ccttatgacc	tgggctacac	acgtgctaca	atggacggta	caaagagctg	1260
caagaccgcg	aggtggagct	aatctcataa	aaccgttctc	agttcrgatt	gtaggctgca	1320
actcgcctac	atgaagctgg	aatcgctagt	aatcgcggat	cagcatgccg	cgggtgaatac	1380
gttcccgggc	cttgtagaca	ccgcccgtca	caccacgaga	gtttgtaaca	cccgaagtcg	1440
gtggggtaac	ctttttggag	ccagccgcct	aaggtgggac	agatgattgg	ggtgaagtcg	1500
taacaaggta	gcc					1513

&lt;210&gt; 6

&lt;211&gt; 1512

&lt;212&gt; DNA

<213> *Bacillus subtilis* ssp. *taxi*

&lt;400&gt; 6

agagtttgat	cctgggtcag	gacgaacgct	ggcggcgtgc	ctaatacatg	caagtcgagc	60
gaacagatgg	gagcttgctc	cctgatgtta	gcggcggacg	ggtgagtaac	acgtgggtaa	120
cctgcctgta	agactgggat	aactccggga	aatcggggct	aataccggat	ggttgtttga	180
accgcatgg	tcagacataa	aaggtggctt	cggctaccac	ttacagatgg	acccgcggcg	240
cattagctag	ttggtgaggt	aacggctcac	caaggcgacg	atgcgtagcc	gacctgagag	300
ggtgatcggc	cacactggga	ctgagacacg	gccagactc	ctacgggagg	cagcagtagg	360
gaatcttccg	caatggacga	aagtctgacg	gagcaacgcc	gcgtgagtga	tgaagggttt	420
cggatcgtaa	agctctgttg	ttaggggaaga	acaagtgccg	ttcaaataag	gcggcacctt	480
gacgggtacct	aaccagaaaag	ccacggctaa	ctacgtgcc	gcagccgcgg	taatacgtag	540
gtggcaagcg	ttgtccggaa	ttattgggcg	taaagggtc	gcaggcggtt	tcttaagtct	600
gatgtgaaag	cccccggttc	aacgggggag	ggtcattgga	aactggggaa	cttgagtgca	660
gaagaggaga	gtggaattcc	acgtgtagcg	gtgaaatgcg	tagagatgtg	gaggaacacc	720
agtggcgaag	gcgactctct	ggtctgtaac	tgacgctgag	gagcgaaagc	gtggggagcg	780
aacaggatta	gataccctgg	tagtccacgc	cgtaaacgat	gagtgtctag	tgttaggggg	840
tttccgcccc	ttagtgctgc	agctaacgca	ttaagcactc	cgcctgggga	gtacgggtcgc	900
aagactgaaa	ctcaaaggaa	ttgacggggg	ccgcacacaag	cgggtggagca	tgtgggtttta	960
ttcgaagcaa	cgcaagaac	cttaccaggt	cttgacatcc	tctgacaatc	ctagagatag	1020
gacgtccctt	tcggggggcag	agtgcacaggt	ggtgcatgg	tgctgctcagc	tcgtgtcgtg	1080
agatgttggg	ttaagtcccc	caacgagcgc	aacccttgat	cttagttgcc	agcattcagt	1140
tgggcactct	aaggtgactg	cgggtgacaa	accggaggaa	ggtgggggatg	acgtcaaatac	1200
atcatgcccc	ttatgacctg	ggctacacac	gtgctacaat	gggcagaaca	aagggcagcg	1260
aaaccgcgag	gttaagccaa	tcccacaaat	ctgttctcag	ttcggatcgc	agtctgcaac	1320
tcgactgcgt	gaagctggaa	tcgctagtaa	tcgcggatca	gcatgccgcg	gtgaatacgt	1380
tcccgggcct	tgtacacacc	gcccgtcaca	ccacgagagt	ttgtaacacc	cgaagtcggt	1440
gaggtaacct	ttatggagcc	agccgccgaa	ggtgggacag	atgattgggg	tgaagtcgta	1500
acaaggtaac	cg					1512

<210> 7  
 <211> 1458  
 <212> DNA  
 <213> Pantoea sp. BCM 2

<400> 7

gcctaacagt	gcgagtcgga	cggtagcaca	gaggagcttg	ctcttcgggt	gacgagtggc	60
ggacgggtga	gtaatgtctg	gggatctgcc	cgatggaggg	ggataaccac	tggaaacggt	120
ggctaatacc	gcataatgtc	gcaagaccaa	agtgggggac	cttcgggcct	cacaccatcg	180
gatgaaccca	gatgggatta	gctagtaggt	ggggtaacgg	ctcacctagg	cgacgatccc	240
tagctggtct	gagaggatga	ccagccacac	tggaaactgag	acacgggtcca	gactcctacg	300
ggaggcagca	gtggggaata	ttgcacaatg	ggcgcaagcc	tgatgcagcc	atgccgcgtg	360
tatgaagaag	gccttcgggt	tgtaaagtac	tttcancggg	gaggaagggg	acgagggttaa	420
taaccccggt	cattgacgtt	acccgcagaa	gaagcaccgg	ctaactccgt	gccagcagcc	480
gcggtaatac	ggaggggtgca	agcggttaatc	ggaattactg	ggcgtaaagc	gcacgcaggc	540
ggtctgttaa	gtcagatgtg	aaatccccgg	gcttaacctg	ggaactgcat	ttgaaactgg	600
caggcttgag	tcttgtagag	gggggtagaa	ttccagggtg	agcggtgaaa	tgcgtagaga	660
tctggaggaa	taccggtggc	gaaggcggcc	ccctggacaa	agactgacgt	tcaggtgcga	720
aagcgtgggg	agcaaacagg	attagatacc	ctggtagtcc	acgccgtaaa	cgatgtcgac	780
ttggaggctg	tgagcatgat	tcgtggcttc	cggagctaac	gcgttaagtc	gaccgcctgg	840
ggagtacggc	cgcaaggtta	aaattcaaata	gaattgacgg	gggcccgcac	aagcggtgga	900
gcatgtggtt	taattcgatg	caacgcgaag	aaccttacct	gytcttgaca	tccacggaat	960
tyggcagaga	tgccttagtg	ccttcgggaa	ccgtgagaca	ggtgctgcat	ggctgtcgtc	1020
agctcgtgtt	gtgaaatgtt	gggttaagtc	ccgcaacgag	cgcaaccctt	atcctttgtt	1080
gccagcgatt	cggtcgggaa	ctcaaaggag	actgccgggtg	ataaacggga	ggaagggtggg	1140
gatgacgtca	agtcatcatg	gcccttacga	gcagggtctac	acacgtgcta	caatggcgca	1200
tacaaagaga	agcgacctcg	cgagagcaag	cggacctcac	aaagtgcgtc	gtagtccgga	1260
tcggagtctg	caactcgact	ccgtgaagtc	ggaatcgcta	gtaatcgtgg	atcagaatgc	1320
cacgggtgaat	acgttcccgg	gccttgtaga	caccgcccgt	cacaccatgg	gagtgggttg	1380
caaaagaagt	aggtagctta	accttcggga	gggcgcttac	cactttgtga	ttcatgactg	1440
gggtgaagtc	gtaacaag					1458

<210> 8  
 <211> 1451  
 <212> DNA  
 <213> Pantoea sp. BCM 3

<400> 8

agtgcgagtc	ggacggtagc	acagagagct	tgctcttggg	tgacgagtgg	cggacgggtg	60
agtaatgtct	ggggatctgc	ccgatagagg	gggataacca	ctggaaacgg	tggctaatac	120
cgcataacgt	cgcaagacca	aagaggggga	ccttcggggc	tctcactatc	ggatgaaccc	180
agatgggatt	agctagtagg	cggggtaatg	gcccacctag	gcgacgatcc	ctagctggtc	240
tgagaggatg	accagccaca	ctggaactga	gacacgggtcc	agactcctac	gggaggcagc	300
agtggggaat	attgcacaat	gggcgcaagc	ctgatgcagc	catgccgcgt	gtatgaagaa	360
ggccttcggg	ttgtaaagta	ctttcagcgg	ggaggaaggc	gatgcggtta	ataaccctgt	420
cgattgacgt	tacccgcaga	agaagcaccg	gctaactccg	tgccagcagc	cgcggttaata	480
cggaggggtg	aagcgttaat	cggaattact	gggcgtaaag	cgcacgcagg	cggctctgtta	540
agtcagatgt	gaaatccccg	ggcttaacct	gggaactgca	tttgaaactg	gcaggcttga	600
gtctttaga	rgggggtaga	attccagggtg	tagcggtgaa	atgcgtagag	atctggagga	660
ataccgggtg	cgaaggcggc	cccctggaca	aagactgacg	ctcagggtgcg	aaagcgtggg	720
gagcaaacag	gattagatac	cctggtagtc	cacgccgtaa	acgatgtcga	cttgagggtt	780
gttcccttga	ggagtggctt	ccggagctaa	cgcgttaagt	cgaccgcctg	gggagtacgg	840
ccgcaagggt	aaaattcaaa	tgaattgacg	ggggcccgcg	caagcgggtg	agcatgtggt	900
ttaattcgat	gcaacgcgaa	gaaccttacc	tactcttgac	atccacggaa	tttyggcagag	960
atgctttggt	gccttcggga	accctgagac	aggtgctgca	tggctgtcgt	cagctcgtgt	1020
tgtgaaatgt	tgggttaagt	ccgcgaacga	gcgcaaccct	tatcctttgt	tgccagcgat	1080



tcggtcggga	actcaaagga	gactgccggt	gataaaccgg	aggaagggtg	ggatgacgtc	1140
aagtcacat	ggcccttacg	agtagggcta	cacacgtgct	acaatggcgc	atacaaagag	1200
aagcgacctc	gcgagagcaa	gcggacctca	caaagtgcgt	cgtagtccgg	atcggagtct	1260
gcaactcgac	tccgtgaagt	cggaaatcgct	agtaatcgct	gatcagaatg	ccacgggtgaa	1320
tacgttcccc	ggccttgtac	acaccgcccc	tcacaccatg	ggagtggggt	gcaaaagaag	1380
taggtagctt	aaccttcggg	agggcgctta	ccactttgtg	attcatgact	ggggtgaagt	1440
cgtaacaagt	a					1451

&lt;210&gt; 9

&lt;211&gt; 1513

&lt;212&gt; DNA

&lt;213&gt; Bacillus megaterium ssp. BCM 9

&lt;400&gt; 9

agagtttgat	catggctcag	gatgaacgct	ggcggcggtgc	ctaatacatg	caagtcgagc	60
gaactgatta	gaagcttgct	tctatgacgt	tagcggcgga	cgggtgagta	acacgtgggc	120
aacctgcctg	taagactggg	ataacttcgg	gaaaccgaag	ctaataccgg	ataggatctt	180
ctccttcacg	ggagatgatt	gaaagatggt	ttcggtctatc	acttacagat	gggcccgcgg	240
tgcattagct	agttgggtgag	gtaacggccc	accaaggcaa	cgatgcatag	ccgacctgag	300
agggatgatc	gccacactgg	gactgagaca	cggcccagac	tcctacggga	ggcagcagta	360
gggaatcttc	cgcaatggac	gaaagtctga	cggagcaacg	ccgcgtgagt	gatgaaggct	420
ttcgggtcgt	aaaactctgt	tgttagggaa	gaacaagtac	gagagtaact	gctcgtacct	480
tgacggtacc	taaccagaaa	gccacggcta	actacgtgcc	agcagccgcg	gtaatacgtg	540
ggtggcaagc	gttatccgga	attattgggc	gtaaagcgcg	ccgcaggcgg	tttcttaagt	600
ctgatgtgaa	agcccacggc	tcaaccgtgg	garggtcatt	ggaaactggg	gaacttgagt	660
gcagaagaga	aaagcggaat	tccacgtgta	gcggtgaaat	gcgtagagat	gtggagggaac	720
accagtggcg	aaggcggtt	tttggtctgt	aactgacgct	gaggcgcgaa	agcgtgggga	780
gcaaacagga	ttagataccc	tggtagtcca	cgcgtaaac	gatgagtgtc	aagtgttaga	840
gggtttccgc	ccttttagtgc	tgcagctaac	gcattaagca	ctccgcctgg	ggagtacggt	900
cgcaagactg	aaactcaaag	gaattgacgg	gggcccgcac	aagcgggtgga	gcatgtgggt	960
taattcgaag	caacgcgaag	aaccttacca	ggtcttgaca	tcctctgaca	actctagaga	1020
tagagctttc	cccttcgggg	gacagagtga	caggtgggtgc	atgggtgtcg	tcagctcgtg	1080
tcgtgagatg	ttgggttaag	tcccgcacac	agcgcacacc	ttgatcttag	ttgccagcat	1140
ttagtggggc	actctaaggt	gactgccggt	gacaaaccgg	aggaagggtg	ggatgacgtc	1200
aaatcatcat	ggcccttatg	acctgggcta	cacacgtgct	acaatggatg	gtacaaaggg	1260
ctgcaagacc	gcgaggtcaa	gccaatccca	taaaaccatt	ctcagttcgg	attgtaggct	1320
gcaactcgcc	tacatgaagc	tggaatcgct	agtaatcgcg	gatcagcatg	ccgcgggtgaa	1380
tacgttcccc	ggccttgtac	acaccgcccc	tcacaccacg	agagtttgta	acaccgaag	1440
tcggtggagt	aaccgtaagg	agctagccgc	ctaagggtggg	acagatgatt	ggggtgaagt	1500
cgtaacaagg	tag					1513

&lt;210&gt; 10

&lt;211&gt; 1452

&lt;212&gt; DNA

&lt;213&gt; Curtobacterium sp. BCM 5

&lt;400&gt; 10

cgctgggcgg	cctgcttaac	acntgcaagt	cgaacgatga	tgcccagctt	gctgggtgga	60
ttagtggcga	acgggtgagt	aacacgtgag	taacctgccc	ctgactctgg	gataagcgtt	120
ggaaacgacg	tctaatactg	gatatgatca	ctggccgcac	ggtctggtgg	tggaagatt	180
ttttgggttg	ggatggactc	gcggcctatc	agcttggttg	tgaggtaatg	gctcaccaag	240
gcgacgacgg	gtagccggcc	tgagaggggtg	accggccaca	ctgggactga	gacacggccc	300
agactcctac	gggaggcaac	agtggggaat	attgcacaat	gggcgaaagc	ctgatgcagc	360
aacgccgcgt	gagggatgac	ggccttcggg	ttgtaaacct	cttttagtag	ggaagaagcg	420
aaagtacggg	tacctgcaga	aaaagcaccg	gctaactacg	tgccagcagc	cgcggttaata	480

cgtagggtgc	aagcgttgct	cggaattatt	gggcgtaaag	agctcgtagg	cggtttgctg	540
cgtctgctgt	gaaatcccga	ggctcaacct	cgggcttgca	gtgggtacgg	gcagactaga	600
gtgcggtagg	ggagattgga	attcctgggtg	tagcgggtgga	atgcgcagat	accaggagga	660
acaccgatgg	cgaaggcaga	tctctggggc	gtaactgacg	ctgaggagcg	aaagcatggg	720
gagcgaacag	gattagatac	cctggtagtc	catgccgtaa	acgttggggc	ctagatgtag	780
ggacctttcc	acggttttctg	tgctcgtagct	aacgcattaa	gcgccccgcc	tggggagtac	840
ggccgcaagg	ctaaaactca	aaggaattga	cggggggccc	cacaagcggc	ggagcatgcg	900
gattaattcg	atgcaacgcg	aagaacctta	ccaaggmttg	acatacaccg	gaaacggcca	960
gagatggctg	cccccttggtg	gtcgggtgtac	aggtgggtgca	tggttgtcgt	cagctcgtgt	1020
cgtgagatgt	tgggttaagt	cccgcacaga	gcgcaaccct	cgttctatgt	tgccagcggg	1080
ttatgccggg	gactcatagg	agactgccgg	ggtcaactcg	gaggaagggtg	gggatgacgt	1140
caaatcatca	tgcccccttat	gtcttggggt	tcacgcacgc	tacaatggcc	ggtacaaagg	1200
gctgcgatac	cgtaagggtgg	agcgaatccc	aaaaagccgg	tctcagttcg	gattgaggtc	1260
tgcaactcga	cctcatgaag	tcggagtcgc	tagtaatcgc	agatcagcaa	cgctgcgggtg	1320
aatacgttcc	cgggccttgt	acacaccgcc	cgtcaagtca	tgaaagtcgg	taacaccgga	1380
agccggtggc	ctaacccttg	tggaaggagc	cgtcgaagggt	gggatcggtg	attaggacta	1440
agtcgtaaca	ag					1452

&lt;210&gt; 11

&lt;211&gt; 1450

&lt;212&gt; DNA

&lt;213&gt; Sphingomonas sp. BCM 7

&lt;400&gt; 11

agagtttgat	catggctcag	aacgaacgct	ggcggcatgc	ctaacacatg	caagtcgaac	60
gagatcttcg	gatctagtgg	cgcacgggtg	cgtaacgcgt	gggaatctgc	cctttgggtc	120
ggaataacag	ttggaaacga	ctgctaata	ccggatgatg	acgtaagtc	aaagatttat	180
cgccagagga	tkagcccgcg	taggattagc	tagttgggtg	ggtaagagcg	caccaaggcg	240
acgatcctta	gctgggtctga	gargatgac	agccacactg	ggactgagac	acggcccaga	300
ctcctacggg	aggcagcagt	ggggaatwtt	ggacaatggg	cgaaagcctg	atccagcaat	360
gccgcgtgag	tgatgaaggc	cttaggggtg	taaagctctt	ttaccgggga	tgataatgac	420
agtaccggga	gaataagctc	cggctaactc	cgtgccagca	gccgcggtaa	tacggaggga	480
gctagcgttg	ttcggaatta	ctgggcgtaa	agcgcacgta	rgcggctttg	taagttagag	540
gtgaaagcct	ggagctcaac	tccagaattg	cctttaagac	tgcatcgctt	gaatccagga	600
gaggtgagtg	gaattccgag	tgtagagggtg	aaattcgtag	atattcggaa	gaacaccagt	660
ggcgaaggcg	gctcactgga	ctgggtattga	cgctgagggtg	cgaaagcgtg	gggagcaaac	720
aggattagat	accctggtag	tccacgccgt	aaacgatgat	aactagctgt	ccggggactt	780
ggtctttggg	tggcgcagct	aacgcattaa	gttatccgcc	tggggagtac	ggccgcaagg	840
ttaaaactca	aatgaattga	cgggggcctg	cacaagcggg	ggagcatgtg	gtttaattcg	900
aagcaacgcg	cagaacctta	ccagcgtttg	acatgtccgg	acgatttctg	gagacagatc	960
tcttcccttc	ggggactgga	acacagggtg	tgcatggctg	tcgtcagctc	gtgtcgtgag	1020
atgttgggtt	aagtcccgcg	acgagcgcaa	ccctcgccct	tagttaccat	catttagttg	1080
gggactctaa	aggaaccgcc	ggtgataagc	cggaggaagg	tggggatgac	gtcaagtcct	1140
catggccctt	acgcgctggg	ctacacacgt	gctacaatgg	cggtgacagt	gggcagcaaa	1200
ctcgcgagag	tgcgctaata	tccaaaagcc	gtctcagttc	ggattgttct	ctgcaactcg	1260
agagcatgaa	ggcggaatcg	ctagtaatcg	cggatcagca	tgccgcgggtg	aatacgttcc	1320
caggccttgt	acacaccgcc	cgtcacacca	tgggagttgg	gttcacccga	aggcgttgcg	1380
ctaactcagc	aatgagaggc	aggcgaccac	ggtgggctta	gcgactgggg	tgaagtcgta	1440
acaaggtaac						1450

&lt;210&gt; 12

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

<400> 12  
agagtttgat cmtggctcag

20

<210> 13  
<211> 22  
<212> DNA  
<213> Artificial Sequence

<400> 13  
tacggytacc ttggttacgac tt

22

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/CA 98/01150

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12P17/02 C12N3/00 C12N1/00 C12N15/01 C12N1/20  
C12P7/62 C12P7/26 //(C12P17/02,C12R1:07),(C12P17/02,  
C12R1:125)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12P C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 04154 A (NIPPON STEEL CORPORATION) 9 February 1995 see page 5; claims 1-10 ---	1-57
X	WO 96 34972 A (BCM DEVELOPMENT INC.) 7 November 1996 cited in the application see the whole document ---	1-57
X	EP 0 668 360 A (BRISTOL-MYERS SQUIBB COMPANY) 23 August 1995 see page 6 - page 7; claims 1-21 --- -/--	1-57

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance  
"E" earlier document but published on or after the international filing date  
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)  
"O" document referring to an oral disclosure, use, exhibition or other means  
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention  
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone  
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.  
"&" document member of the same patent family

Date of the actual completion of the international search

27 May 1999

Date of mailing of the international search report

10/06/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Douschan, K



# INTERNATIONAL SEARCH REPORT

Int lional Application No  
PCT/CA 98/01150

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, L	CHEMICAL ABSTRACTS, vol. 130, 1999 Columbus, Ohio, US; abstract no. 174870, JAZIRI, M. ET AL: "A new concept for the isolation of bioactive taxoids" XP002103986	1-40
Y	see abstract & CURR. TOP. PHYTOCHEM. (1997), 1, 67-76 CODEN: CTPHFK,	41-57
X	--- CHEMICAL ABSTRACTS, vol. 128, Columbus, Ohio, US; abstract no. 202215, HANSON, RONALD L. ET AL: "Enzymic hydrolysis of 7-xylosyltaxanes by xylosidase from Moraxella sp." XP002103987	6
Y	see abstract & BIOTECHNOL. APPL. BIOCHEM. (1997), 26(3), 153-158 CODEN: BABIEC; ISSN: 0885-4513,	1-5, 7-57
X	--- WO 96 32490 A (NOVOPHARM LIMITED) 17 October 1996	6
Y	see page 1 - page 6; claims 1-9	1-5, 7-57
X	--- WO 97 16200 A (MERCK & CO., INC.) 9 May 1997 cited in the application	6
Y	see the whole document	18-30
Y	--- WO 93 21338 A (THE RESEARCH AND DEVELOPMENT INSTITUTE, INC.) 28 October 1993 cited in the application see claims 1-33	1-57
Y	--- STROBEL, G. ET AL.: "Taxol from Pestalotiopsis microspora, an endophytic fungus of Taxus wallachiana" MICROBIOLOGY, vol. 142, 1996, pages 435-440, XP002103985 cited in the application see the whole document -----	1-57

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 98/01150

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9504154 A	09-02-1995	JP 7177879 A	18-07-1995
WO 9634972 A	07-11-1996	US 5561055 A	01-10-1996
		AU 5494696 A	21-11-1996
		CA 2220081 A	07-11-1996
EP 0668360 A	23-08-1995	AT 178945 T	15-04-1999
		AU 693827 B	09-07-1998
		AU 1015695 A	27-07-1995
		CA 2138674 A	14-07-1995
		CN 1111229 A	08-11-1995
		DE 69508987 D	20-05-1999
		FI 950137 A	14-07-1995
		HU 70987 A	28-11-1995
		JP 7203981 A	08-08-1995
		US 5700669 A	23-12-1997
WO 9632490 A	17-10-1996	AU 5265696 A	30-10-1996
WO 9716200 A	09-05-1997	AU 7529396 A	22-05-1997
WO 9321338 A	28-10-1993	US 5322779 A	21-06-1994
		AU 675428 B	06-02-1997
		AU 4102093 A	18-11-1993
		CA 2140935 A	17-10-1993
		EP 0672168 A	20-09-1995
		JP 7505778 T	29-06-1995
		US 5861302 A	19-01-1994

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**